

q -SERIES AND TAILS OF COLORED JONES POLYNOMIALS

PAUL BEIRNE AND ROBERT OSBURN

ABSTRACT. We extend the table of Garoufalidis, Lê and Zagier concerning conjectural Rogers-Ramanujan type identities for tails of colored Jones polynomials to all alternating knots up to 10 crossings. We then prove these new identities using q -series techniques.

1. INTRODUCTION

The colored Jones polynomial $J_N(K; q)$ for a knot K is an important quantum invariant of 3-manifolds. Here, we use the normalization $J_1(K; q) = 1$ for all knots K and $J_2(K; q)$ is the Jones polynomial of K . The *tail* of $J_N(K; q)$ is a power series whose first N coefficients agree (up to a common sign) with the first N coefficients for $J_N(K; q)$ for all $N \geq 1$. If K is an alternating knot, then the tail exists and equals an explicit q -multisum $\Phi_K(q)$ (see [1], [3], [5]).

Recently, Garoufalidis and Lê (with Zagier) presented a table (see Table 6 in [5]) of 43 conjectural Rogers-Ramanujan type identities between the tails $\Phi_K(q)$ and products of theta functions and/or false theta functions. This table consisted of the following knots K : all alternating knots up to 8_4 , the twist knots K_p , $p > 0$ or $p < 0$, the torus knots $T(2, p)$, $p > 0$, each of their mirror knots $-K$ and -8_5 . For example, if we define for a positive integer b

$$h_b = h_b(q) = \sum_{n \in \mathbb{Z}} \epsilon_b(n) q^{\frac{bn(n+1)}{2} - n}$$

where

$$\epsilon_b(n) = \begin{cases} (-1)^n & \text{if } b \text{ is odd,} \\ 1 & \text{if } b \text{ is even and } n \geq 0, \\ -1 & \text{if } b \text{ is even and } n < 0 \end{cases}$$

and

$$(a)_n = (a; q)_n = \prod_{k=1}^n (1 - aq^{k-1}),$$

valid for $n \in \mathbb{N} \cup \{\infty\}$, then

Date: January 21, 2016.

2010 *Mathematics Subject Classification.* Primary: 33D15; Secondary: 05A30, 57M25.

Key words and phrases. q -series identities, colored Jones polynomial, tails.

$$\begin{aligned} \Phi_{7_2}(q) &= (q)_\infty^7 \sum_{a,b,c,d,e,f,g \geq 0} \frac{q^{3a^2+2a+b^2+bg+ac+ad+ae+af+ag+cd+de+ef+fg+c+d+e+f+g}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_{b+g}(q)_{a+c}(q)_{a+d}(q)_{a+e}(q)_{a+f}(q)_{a+g}} \\ &\stackrel{?}{=} h_6. \end{aligned} \tag{1.1}$$

Note that $h_1(q) = 0$, $h_2(q) = 1$ and $h_3(q) = (q)_\infty$. In general, h_b is a theta function if b is odd and a false theta function if b is even. Using q -series techniques, Keilthy and the second author [8] proved not only (1.1), but all of the remaining conjectural identities in [5].

A natural question (due to Zagier) is whether Table 6 in [5] is “complete”, i.e., if we consider an alternating knot K not in this table, then does the evaluation of $\Phi_K(q)$ already appear? The purpose of this paper is to demonstrate that the answer is no. In particular, we first extend the table of Garoufalidis, Lê and Zagier to include all alternating knots up to 10 crossings. Our second goal is to prove the following main result.

Theorem 1.1. *The identities in Tables 1 and 2 are true.*

K	$\Phi_K(q)$	$\Phi_{-K}(q)$	K	$\Phi_K(q)$	$\Phi_{-K}(q)$	K	$\Phi_K(q)$	$\Phi_{-K}(q)$
8 ₆	h_3h_4	h_5	9 ₆	h_3h_6	h_4	9 ₂₄	h_3h_4	?
8 ₇	h_3h_5	h_3^2	9 ₇	h_3h_4	h_6	9 ₂₅	h_3^3	?
8 ₈	h_3h_5	h_3^2	9 ₈	h_3h_6	h_3^2	9 ₂₆	$h_3^2h_4$	h_3^3
8 ₉	h_3h_4	h_3h_4	9 ₉	h_4h_5	h_4	9 ₂₇	h_3^3	$h_3^2h_4$
8 ₁₀	?	h_3^2	9 ₁₀	h_4^2	h_5	9 ₂₈	?	?
8 ₁₁	h_3h_4	h_3h_4	9 ₁₁	h_4h_5	h_3^2	9 ₂₉	?	?
8 ₁₂	h_3h_4	h_3h_4	9 ₁₂	h_3h_4	h_3h_5	9 ₃₀	h_3^3	?
8 ₁₃	$h_3^2h_4$	h_3^2	9 ₁₃	h_4^2	h_3h_4	9 ₃₁	h_3^4	h_3^3
8 ₁₄	h_3h_4	h_3^3	9 ₁₄	$h_3^2h_5$	h_3^2	9 ₃₂	?	?
8 ₁₅	h_3^3	?	9 ₁₅	h_3h_4	h_3h_5	9 ₃₃	?	?
8 ₁₆	?	?	9 ₁₆	h_4	?	9 ₃₄	?	?
8 ₁₇	?	?	9 ₁₇	h_3^2	$h_3^2h_5$	9 ₃₅	?	h_3
8 ₁₈	?	?	9 ₁₈	h_3h_4	h_4^2	9 ₃₆	?	h_3^2
9 ₁	h_9	1	9 ₁₉	h_3h_5	h_3^3	9 ₃₇	h_3^3	?
9 ₂	h_8	h_3	9 ₂₀	h_3^2	$h_3h_4^2$	9 ₃₈	?	?
9 ₃	h_7	h_4	9 ₂₁	h_3h_4	$h_3^2h_4$	9 ₃₉	?	?
9 ₄	h_6	h_5	9 ₂₂	?	h_3^2	9 ₄₀	?	?
9 ₅	h_3	h_4h_6	9 ₂₃	h_4^2	h_3^3	9 ₄₁	?	?

TABLE 1.

Unfortunately, we were unable to find similar identities not only in each case labelled “?” in Tables 1 and 2, but for any alternating knot (or its mirror) from 10₇₈ to 10₁₂₃. This is also the situation for 8₅ where although one has (after simplification)

$$\Phi_{8_5}(q) = (q)_\infty^2 \sum_{a,b \geq 0} \frac{q^{a^2+a+b^2+b}(q)_{a+b}}{(q)_a^2(q)_b^2}, \tag{1.2}$$

K	$\Phi_K(q)$	$\Phi_{-K}(q)$	K	$\Phi_K(q)$	$\Phi_{-K}(q)$	K	$\Phi_K(q)$	$\Phi_{-K}(q)$
10 ₁	h_9	h_3	10 ₂₆	$h_3h_4^2$	h_3h_4	10 ₅₂	?	h_3^3
10 ₂	?	h_3	10 ₂₇	h_3h_5	$h_3^2h_4$	10 ₅₃	?	h_3^3
10 ₃	h_7	h_5	10 ₂₈	$h_3h_4h_5$	h_3^2	10 ₅₄	?	h_3^2
10 ₄	?	h_3	10 ₂₉	$h_3h_4^2$	h_3h_4	10 ₅₅	?	h_3^3
10 ₅	h_3h_7	h_3^2	10 ₃₀	$h_3h_4^2$	h_3^3	10 ₅₆	?	h_3h_4
10 ₆	h_3h_6	h_5	10 ₃₁	h_3h_5	$h_3^2h_4$	10 ₅₇	?	$h_3^2h_4$
10 ₇	h_3h_6	h_3h_4	10 ₃₂	$h_3h_4^2$	h_3^3	10 ₅₈	?	h_3^3
10 ₈	h_3	h_5h_6	10 ₃₃	$h_3^2h_4$	$h_3^2h_4$	10 ₅₉	?	h_3^3
10 ₉	h_3h_6	h_3h_4	10 ₃₄	h_3h_7	h_3^2	10 ₆₀	?	h_3^3
10 ₁₀	$h_3^2h_6$	h_3^2	10 ₃₅	h_3h_6	h_3h_4	10 ₆₁	?	h_3
10 ₁₁	h_4h_5	h_5	10 ₃₆	h_3h_6	h_3^3	10 ₆₂	?	h_3^2
10 ₁₂	h_3h_5	h_3h_5	10 ₃₇	h_3h_5	h_3h_5	10 ₆₃	?	h_3h_4
10 ₁₃	h_4h_5	h_3h_4	10 ₃₈	h_4h_5	h_3^3	10 ₆₄	?	h_3h_4
10 ₁₄	$h_3^2h_5$	h_3h_4	10 ₃₉	h_3h_4	$h_3^2h_5$	10 ₆₅	?	$h_3^2h_4$
10 ₁₅	h_5^2	h_3^2	10 ₄₀	$h_3^2h_4$	$h_3^2h_4$	10 ₆₆	?	$h_3^2h_5$
10 ₁₆	h_4h_5	h_3h_4	10 ₄₁	$h_3h_4^2$	h_3^3	10 ₆₇	?	h_3^3
10 ₁₇	h_3h_5	h_3h_5	10 ₄₂	$h_3^2h_4$?	10 ₆₈	?	h_3^2
10 ₁₈	$h_3^2h_5$	h_3h_4	10 ₄₃	$h_3^2h_4$	$h_3^2h_4$	10 ₆₉	?	?
10 ₁₉	$h_3h_4h_5$	h_3^2	10 ₄₄	$h_3^3h_4$	h_3^4	10 ₇₁	?	$h_3^2h_4$
10 ₂₀	h_7	h_3h_4	10 ₄₆	?	h_3	10 ₇₂	h_3h_4	?
10 ₂₁	h_3h_6	h_3h_4	10 ₄₇	?	h_3^2	10 ₇₃	?	$h_3^2h_4$
10 ₂₂	h_3h_4	h_4h_5	10 ₄₈	?	h_3h_5	10 ₇₄	?	h_3h_4
10 ₂₃	h_3h_5	$h_3^2h_4$	10 ₄₉	?	$h_3^2h_5$	10 ₇₅	?	?
10 ₂₄	h_4h_5	h_3h_4	10 ₅₀	?	h_3h_4	10 ₇₆	?	h_5
10 ₂₅	$h_3h_4^2$	h_3h_4	10 ₅₁	?	$h_3^2h_4$	10 ₇₇	?	h_3h_5

TABLE 2.

the modular (or false theta, mock/mixed mock, quantum modular) properties of the double sum in (1.2) are not clear. Another approach to Theorem 1.1 is to utilize the skein-theoretic techniques in [2], [4] and [7]. It would be of considerable interest to investigate the connection between skein theory and *q*-series to gain a better understanding of these unknown cases and of a general framework.

Finally, it would be desirable to study *q*-series identities in other settings which arise from knot theory. For example, the *q*-multisum $\Phi_K(q)$ occurs as the “0-limit” of $J_N(K; q)$ (see Theorem 2 in [5]). Garoufalidis and Lê have also obtained an explicit formula (see Theorem 3 in [5]) for the “1-limit” of $J_N(K; q)$. Also, do tails exist for two generalizations of $J_N(K; q)$, namely for colored HOMFLY polynomials and colored superpolynomials (see [6], [9]–[11])?

The paper is organized as follows. In Section 2, we recall the necessary background from [8]. In Section 3, we prove Theorem 1.1.

2. PRELIMINARIES

We first recall six *q*-series identities (see (2.1)–(2.3), Lemma 2.1, (4.3) and the proof of (4.1) in [8]). Namely,

$$\sum_{n=0}^{\infty} \frac{t^n}{(q)_n} = \frac{1}{(t)_{\infty}}, \quad (2.1)$$

$$\sum_{n=0}^{\infty} \frac{(-1)^n t^n q^{n(n-1)/2}}{(q)_n} = (t)_{\infty}, \quad (2.2)$$

$$\sum_{n=0}^{\infty} \frac{q^{n^2+An}}{(q)_n(q)_{n+A}} = \frac{1}{(q)_{\infty}} \quad (2.3)$$

for any integer A ,

$$\sum_{m,n \geq 0} (-1)^n \frac{q^{m^2+m+mn+\frac{n(n+1)}{2}}}{(q)_m(q)_n} = h_4, \quad (2.4)$$

$$\sum_{l,m,n \geq 0} (-1)^{l+n} \frac{q^{\frac{3l(l+1)}{2}+m^2+m+\frac{n(n+1)}{2}+2lm+ln+mn}}{(q)_l(q)_m(q)_n} = h_5 \quad (2.5)$$

and

$$\sum_{a \geq 0} (-1)^{na} \frac{q^{\frac{na(a+1)}{2}-a+a \sum_{k=1}^{n-1} c_k}}{(q)_a \prod_{k=1}^{n-1} (q)_{a+c_k}} = \frac{1}{(q)_{\infty}} \sum_{i_1, \dots, i_{n-2} \geq 0} (-1)^{\sum_{k=1}^{n-2} \sum_{j=1}^k i_j} q^{\frac{1}{2} \sum_{k=1}^{n-2} \left(\sum_{j=1}^k i_j \right) \left(1 + \sum_{j=1}^k i_j \right) + \sum_{k=2}^{n-1} \sum_{j=1}^{k-1} c_k i_j} \frac{\prod_{k=1}^{n-2} (q)_{i_k} \prod_{k=1}^{n-2} (q)}{\prod_{k=1}^{n-2} (q)_{c_k + \sum_{j=1}^k i_j}} \quad (2.6)$$

for any $n > 2$ and integers c_k .

Let K be an alternating knot with c crossings and \mathcal{T}_K its associated Tait graph. The reduced Tait graph \mathcal{T}'_K is obtained from \mathcal{T}_K by replacing every set of two edges that connect the same two vertices by a single edge. The tail $\Phi_K(q)$ is given by

$$\Phi_K(q) = (q)_{\infty}^c S_K(q) \quad (2.7)$$

where $S_K(q)$ is an explicitly constructed q -multisum (see pages 261–264 in [8]). Now, by Theorem 2 in [2] or Corollary 1.12 in [5], if $\mathcal{T}'_K \cong \mathcal{T}'_L$ for two alternating knots K and L , then $\Phi_K(q) = \Phi_L(q)$. Thus, by comparing the reduced Tait graphs for those knots in Table 1 of [8] and Tables 1 and 2 above, it suffices to verify the conjectural identities in the following cases: $8_7, 8_{13}, -9_5, 9_{14}, -9_{17}, -9_{20}, -9_{27}, 9_{31}, 10_5, -10_8, 10_{10}, 10_{15}, 10_{19}, 10_{26}, 10_{28}, 10_{44}$.

The strategy for proving Theorem 1.1 is now as follows. For each of the 16 cases, we first compute $S_K(q)$ using the methods from [8]. We then employ (2.1)–(2.6) to reduce this q -multisum to (1.1) or one of the following key identities proven in [8]:

$$S_{51}(q) := \sum_{a,b,c,d,e \geq 0} (-1)^a \frac{q^{\frac{a(5a+3)}{2}+ab+ac+ad+ae+bc+cd+de+b+c+d+e}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_{a+b}(q)_{a+c}(q)_{a+d}(q)_{a+e}} = \frac{1}{(q)_{\infty}^5} h_5, \quad (2.8)$$

$$S_{6_2}(q) := \sum_{a,b,c,d,e,f \geq 0} (-1)^e \frac{q^{2f^2+f+\frac{e(3e+1)}{2}+ab+af+bc+bf+cd+ce+cf+de+a+b+c+d}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_{a+f}(q)_{b+f}(q)_{c+e}(q)_{c+f}(q)_{d+e}} = \frac{1}{(q)_\infty^5} h_4, \quad (2.9)$$

$$\begin{aligned} S_{7_1}(q) &:= \sum_{a,b,c,d,e,f,g \geq 0} (-1)^a \frac{q^{\frac{a(7a+5)}{2}+ab+ac+ad+ae+af+ag+bc+cd+de+ef+fg+b+c+d+e+f+g}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_{a+b}(q)_{a+c}(q)_{a+d}(q)_{a+e}(q)_{a+f}(q)_{a+g}} \\ &= \frac{1}{(q)_\infty^7} h_7, \end{aligned} \quad (2.10)$$

$$\begin{aligned} S_{7_4}(q) &:= \sum_{a,b,c,d,e,f,g \geq 0} \frac{q^{2f^2+f+2g^2+g+ab+ag+bc+bg+cd+cf+cg+de+df+ef+a+b+c+d+e}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_{a+g}(q)_{b+g}(q)_{c+f}(q)_{c+g}(q)_{d+f}(q)_{e+f}} \\ &= \frac{1}{(q)_\infty^7} h_4^2, \end{aligned} \quad (2.11)$$

$$\begin{aligned} S_{7_7}(q) &:= \sum_{a,b,c,d,e,f,g \geq 0} (-1)^{e+f+g} \frac{q^{\frac{3a^2}{2}+\frac{e}{2}+\frac{3f^2}{2}+\frac{f}{2}+\frac{3g^2}{2}+\frac{g}{2}+ab+ad+ae+af+bf+cd+cg+de+dg+a+b+c}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_{a+e}(q)_{d+e}(q)_{a+f}(q)_{b+f}(q)_{c+g}} \\ &\times \frac{q^d}{(q)_{d+g}} \\ &= \frac{1}{(q)_\infty^4}, \end{aligned} \quad (2.12)$$

$$\begin{aligned} S_{8_2}(q) &:= \sum_{a,b,c,d,e,f,g,h \geq 0} (-1)^b \frac{q^{3a^2+2a+\frac{b(3b+1)}{2}+ad+ae+af+ag+ah+bc+bd+cd+de+ef+fg+gh+c+d+e+f}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_{b+c}(q)_{b+d}(q)_{a+d}(q)_{a+e}(q)_{a+f}} \\ &\times \frac{q^{g+h}}{(q)_{a+g}(q)_{a+h}} \\ &= \frac{1}{(q)_\infty^7} h_6 \end{aligned} \quad (2.13)$$

and

$$\begin{aligned}
S_{-8_4}(q) &:= \sum_{a,b,c,d,e,f,g,h \geq 0} (-1)^g \frac{q^{\frac{g(5g+3)}{2} + h(2h+1) + ab + ah + bc + bh + cd + cg + ch + de + dg + ef + eg + fg + a + b + c + d}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_{a+h}(q)_{b+h}(q)_{c+g}(q)_{c+h}(q)_{d+g}} \\
&\times \frac{q^{e+f}}{(q)_{e+g}(q)_{f+g}} \\
&= \frac{1}{(q)_\infty^8} h_4 h_5.
\end{aligned} \tag{2.14}$$

3. PROOF OF THEOREM 1.1

Proof of Theorem 1.1. We give full details for 8_7 , -9_5 and -10_8 . As the remaining cases are handled similarly, we sketch their proofs. For $\Phi_{8_7}(q)$, it suffices to prove

$$\begin{aligned}
S_{8_7}(q) &:= \sum_{a,b,c,d,e,g,h,i \geq 0} (-1)^{h+i} \frac{q^{\frac{i(5i+3)}{2} + \frac{h(3h+1)}{2} + g^2 + ab + ag + ah + bc + bh + bi + cd + ci + de + di + ei + a + b + c}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_g(q)_h(q)_i(q)_{a+g}(q)_{a+h}(q)_{b+h}(q)_{b+i}(q)_{c+i}} \\
&\times \frac{q^{d+e}}{(q)_{d+i}(q)_{e+i}} \\
&= \frac{1}{(q)_\infty^7} h_5.
\end{aligned} \tag{3.1}$$

We now have

$$\begin{aligned}
S_{8_7}(q) &= \frac{1}{(q)_\infty} \sum_{a,b,c,d,e,h,i \geq 0} (-1)^{h+i} \frac{q^{\frac{i(5i+3)}{2} + \frac{h(3h+1)}{2} + ab + ah + bc + bh + bi + cd + ci + de + di + ei + a + b + c}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_h(q)_i(q)_{a+h}(q)_{b+h}(q)_{b+i}(q)_{c+i}(q)_{d+i}} \\
&\times \frac{q^{d+e}}{(q)_{e+i}} \\
&\text{(evaluate the } g\text{-sum with (2.3))} \\
&= \frac{1}{(q)_\infty^2} \sum_{a,b,c,d,e,h,i \geq 0} (-1)^{h+i} \frac{q^{\frac{i(5i+3)}{2} + \frac{h(h+1)}{2} + ab + ah + bc + bi + cd + ci + de + di + ei + a + b + c + d + e}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_h(q)_i(q)_{b+h}(q)_{b+i}(q)_{c+i}(q)_{d+i}(q)_{e+i}} \\
&\text{(apply (2.6) to the } h\text{-sum with } n = 3\text{)} \\
&= \frac{1}{(q)_\infty^2} \sum_{b,c,d,e,i \geq 0} (-1)^i \frac{q^{\frac{i(5i+3)}{2} + bc + bi + cd + ci + de + di + ei + b + c + d + e}}{(q)_b(q)_c(q)_d(q)_e(q)_i(q)_{b+i}(q)_{c+i}(q)_{d+i}(q)_{e+i}} \\
&\text{(evaluate the } a\text{-sum with (2.1), simplify, then use (2.2) for the } h\text{-sum).}
\end{aligned}$$

Thus, (3.1) then follows from (2.8) after letting $i \rightarrow a$.

For $\Phi_{8_{13}}(q)$, it suffices to prove

$$\begin{aligned}
 S_{8_{13}}(q) &:= \sum_{a,c,d,e,f,g,h,i \geq 0} (-1)^{g+h} \frac{q^{\frac{g(3g+1)}{2} + \frac{(3h+1)}{2} + i(2i+1) + af + ag + ci + cd + de + di + ef + eh + ei}}{(q)_a(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_{a+g}(q)_{c+i}(q)_{d+i}(q)_{e+i}(q)_{e+h}} \\
 &\quad \times \frac{q^{fh+fg+a+c+d+e+f}}{(q)_{f+h}(q)_{f+g}} \\
 &= \frac{1}{(q)_\infty^6} h_4.
 \end{aligned} \tag{3.2}$$

Apply (2.6) with $n = 3$ to the g -sum, (2.1) to the a -sum, then simplify and (2.2) to the g -sum to obtain

$$S_{8_{13}}(q) = \frac{1}{(q)_\infty} \sum_{c,d,e,f,h,i \geq 0} (-1)^h \frac{q^{\frac{h(3h+1)}{2} + i(2i+1) + ci + cd + de + di + ef + eh + ei + fh + c + d + e + f}}{(q)_c(q)_d(q)_e(q)_f(q)_h(q)_i(q)_{c+i}(q)_{d+i}(q)_{e+i}(q)_{e+h}(q)_{f+h}}.$$

Thus, (3.2) then follows from (2.9) upon $(c, d, e, f, h, i) \rightarrow (a, b, c, d, e, f)$.

For $\Phi_{-9_5}(q)$, it suffices to prove

$$\begin{aligned}
 S_{-9_5}(q) &:= \sum_{a,b,c,d,e,f,g,h,j \geq 0} \frac{q^{h(2h+1) + j(3j+2) + ab + ag + ah + aj + bc + bh + ch + de + dj + ef + ej + fg + fj + gj + a + b + c}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_j(q)_{a+h}(q)_{a+j}(q)_{b+h}(q)_{c+h}(q)_{d+j}} \\
 &\quad \times \frac{q^{d+e+f+g}}{(q)_{e+j}(q)_{f+j}(q)_{g+j}} \\
 &= \frac{1}{(q)_\infty^9} h_4 h_6.
 \end{aligned} \tag{3.3}$$

We now have

$$\begin{aligned}
 S_{-9_5}(q) &= \frac{1}{(q)_\infty} \sum_{a,b,c,d,e,f,g,j,s,t \geq 0} \frac{q^{s^2 + s + st + \frac{t(t+1)}{2} + bs + c(s+t) + j(3j+2) + ab + ag + aj + bc + de + dj + ef + ej + fg}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_j(q)_s(q)_t(q)_{a+j}(q)_{d+j}(q)_{s+a}} \\
 &\quad \times \frac{q^{fj + gj + a + b + c + d + e + f + g}}{(q)_{s+t+b}(q)_{e+j}(q)_{f+j}(q)_{g+j}} \\
 &\quad (\text{apply (2.6) to the } h\text{-sum with } n = 4) \\
 &= \frac{1}{(q)_\infty^2} \sum_{a,b,d,e,f,g,j,s,t \geq 0} \frac{q^{s^2 + s + st + \frac{t(t+1)}{2} + bs + j(3j+2) + ab + ag + aj + de + dj + ef + ej + fg + fj + gj + a + b + d + e}}{(q)_a(q)_b(q)_d(q)_e(q)_f(q)_g(q)_j(q)_s(q)_t(q)_{a+j}(q)_{d+j}(q)_{e+j}(q)_{f+j}(q)_{g+j}} \\
 &\quad \times \frac{q^{f+g}}{(q)_{s+a}} \\
 &\quad (\text{evaluate the } c\text{-sum with (2.1) and simplify})
 \end{aligned}$$

$$= \frac{1}{(q)_\infty^3} h_4 \sum_{a,d,e,g,j \geq 0} \frac{q^{j(3j+2)+ag+aj+de+dj+ej+fg+fj+gj+a+d+e+f+g}}{(q)_a(q)_d(q)_e(q)_f(q)_g(q)_j(q)_{a+j}(q)_{d+j}(q)_{e+j}(q)_{f+j}(q)_{g+j}}$$

(evaluate the b -sum with (2.1), simplify, then apply (2.4) to the st -sum).

Now, (3.3) follows from first applying (2.3) the b -sum in (1.1), then letting $(a, d, e, f, g, j) \rightarrow (c, g, f, e, d, a)$.

For $\Phi_{9_{14}}(q)$, it suffices to prove

$$\begin{aligned} S_{9_{14}}(q) &:= \sum_{a,b,c,d,e,g,h,i,j \geq 0} (-1)^{h+i+j} \frac{q^{\frac{h(3h+1)}{2} + \frac{i(3i+1)}{2} + \frac{j(5j+3)}{2} + ab + ag + ah + ai + bc + bi + bj + cd + cj + de + dj + ej}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_g(q)_h(q)_i(q)_j(q)_{a+h}(q)_{a+i}(q)_{b+i}(q)_{b+j}} \\ &\quad \times \frac{q^{gh+a+b+c+d+e+g}}{(q)_{c+j}(q)_{d+j}(q)_{e+j}(q)_{g+h}} \\ &= \frac{1}{(q)_\infty^7} h_5. \end{aligned} \tag{3.4}$$

First, apply (2.6) with $n = 3$ to the h -sum, (2.1) to the g -sum, simplify and (2.2) to the h -sum, then (2.6) with $n = 3$ to the i -sum, (2.1) to the a -sum, simplify and (2.2) to the i -sum to obtain

$$S_{9_{14}}(q) = \frac{1}{(q)_\infty^2} \sum_{b,c,d,e,j} (-1)^j \frac{q^{\frac{j(5j+3)}{2} + bc + bj + cd + cj + de + dj + ej + b + c + d + e}}{(q)_b(q)_c(q)_d(q)_e(q)_j(q)_{b+j}(q)_{c+j}(q)_{d+j}(q)_{e+j}}.$$

Thus, (3.4) follows from (2.8) after $j \rightarrow a$.

For $\Phi_{-9_{17}}(q)$, it suffices to prove

$$\begin{aligned} S_{-9_{17}}(q) &:= \sum_{a,b,c,d,e,f,h,i,j \geq 0} (-1)^{h+i+j} \frac{q^{\frac{h(3h+1)}{2} + \frac{i(5i+3)}{2} + \frac{j(3j+1)}{2} + ab + aj + bc + bi + bj + cd + ci + de + di + ef}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_h(q)_i(q)_j(q)_{a+j}(q)_{b+i}(q)_{b+j}(q)_{c+i}} \\ &\quad \times \frac{q^{eh+ei+fh+a+b+c+d+e+f}}{(q)_{d+i}(q)_{e+h}(q)_{e+i}(q)_{f+h}} \\ &= \frac{1}{(q)_\infty^7} h_5. \end{aligned} \tag{3.5}$$

First, apply (2.6) with $n = 3$ to the h -sum, (2.1) to the f -sum, simplify and (2.3) to the h -sum, then (2.6) with $n = 3$ to the j -sum, (2.1) to the a -sum, simplify and (2.3) to the j -sum to get

$$S_{-9_{17}}(q) = \frac{1}{(q)_\infty^2} \sum_{b,c,d,e,i \geq 0} (-1)^i \frac{q^{\frac{i(5i+3)}{2} + bc + bi + cd + ci + de + di + ei + b + c + d + e}}{(q)_b(q)_c(q)_d(q)_e(q)_i(q)_{b+i}(q)_{c+i}(q)_{d+i}(q)_{e+i}}.$$

Thus, (3.5) follows from (2.8) after $i \rightarrow a$.

For $\Phi_{-9_{20}}(q)$, it suffices to prove

$$\begin{aligned}
 S_{-9_{20}}(q) &:= \sum_{a,b,c,d,e,f,h,i,j \geq 0} (-1)^h \frac{q^{\frac{h(3h+1)}{2} + i(2i+1) + j(2j+1) + ab + ah + bc + bh + bi + cd + ci + de + di + dj + ef + ej}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_h(q)_i(q)_j(q)_{a+h}(q)_{b+h}(q)_{b+i}(q)_{c+i}} \\
 &\times \frac{q^{fj+a+b+c+d+e+f}}{(q)_{d+i}(q)_{d+j}(q)_{e+j}(q)_{f+j}} \\
 &= \frac{1}{(q)_\infty^8} h_4^2.
 \end{aligned} \tag{3.6}$$

Apply (2.6) with $n = 3$ to the h -sum, (2.1) to the a -sum and simplify, then (2.2) to the h -sum to obtain

$$S_{-9_{20}}(q) = \frac{1}{(q)_\infty} \sum_{b,c,d,e,f,i,j \geq 0} \frac{q^{i(2i+1) + j(2j+1) + bc + bi + cd + ci + de + di + dj + ef + ej + fj + b + c + d + e + f}}{(q)_b(q)_c(q)_d(q)_e(q)_f(q)_i(q)_j(q)_{b+i}(q)_{c+i}(q)_{d+i}(q)_{d+j}(q)_{e+j}(q)_{f+j}}.$$

Now, (3.6) follows from (2.11) after the substitution $(b, c, d, e, f, i, j) \rightarrow (a, b, c, d, e, g, f)$.

For $\Phi_{-9_{27}}(q)$, it suffices to prove

$$\begin{aligned}
 S_{-9_{27}}(q) &:= \sum_{a,b,c,d,e,f,g,h,i \geq 0} (-1)^{f+h} \frac{q^{\frac{f(3f+1)}{2} + g(2g+1) + \frac{h(3h+1)}{2} + i^2 + ab + af + bc + bf + bg + cd + cg + de + dg + dh}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_{a+f}(q)_{b+f}(q)_{b+g}(q)_{c+g}} \\
 &\times \frac{q^{eh+ei+a+b+c+d+e}}{(q)_{d+g}(q)_{d+h}(q)_{e+h}(q)_{e+i}} \\
 &= \frac{1}{(q)_\infty^7} h_4.
 \end{aligned} \tag{3.7}$$

Apply (2.3) to the i -sum, (2.6) with $n = 3$ to the f -sum, (2.1) to the a -sum, simplify and (2.2) to the f -sum to obtain

$$S_{-9_{27}} = \frac{1}{(q)_\infty^2} \sum_{b,c,d,e,g,h \geq 0} (-1)^h \frac{q^{g(2g+1) + \frac{h(3h+1)}{2} + bc + bg + cd + cg + de + dg + dh + eh + b + c + d + e}}{(q)_b(q)_c(q)_d(q)_e(q)_g(q)_h(q)_{b+g}(q)_{c+g}(q)_{d+g}(q)_{d+h}(q)_{e+h}}.$$

Now, (3.7) follows from (2.9) after letting $(b, c, d, e, g, h) \rightarrow (a, b, c, d, f, e)$.

For $\Phi_{9_{31}}(q)$, it suffices to prove

$$\begin{aligned}
S_{9_{31}}(q) &:= \sum_{a,b,c,e,f,g,h,i,j \geq 0} (-1)^{g+h+i+j} \frac{q^{\frac{g(3g+1)}{2} + \frac{h(3h+1)}{2} + \frac{i(3i+1)}{2} + \frac{j(3j+1)}{2} + ab + af + ag + aj + bc + bg + bh}}{(q)_a(q)_b(q)_c(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_a+g(q)_a+j(q)_b+g} \\
&\times \frac{q^{ch+ef+ei+fi+fj+a+b+c+e+f}}{(q)_{b+h}(q)_{c+h}(q)_{e+i}(q)_{f+i}(q)_{f+j}} \\
&= \frac{1}{(q)_\infty^5}.
\end{aligned} \tag{3.8}$$

Apply (2.6) with $n = 3$ to the h -sum, (2.1) to the c -sum, simplify and (2.2) to the h -sum to obtain

$$\begin{aligned}
S_{9_{31}}(q) &= \frac{1}{(q)_\infty} \sum_{a,b,e,f,g,i,j \geq 0} (-1)^{g+i+j} \frac{q^{\frac{g(3g+1)}{2} + \frac{i(3i+1)}{2} + \frac{j(3j+1)}{2} + ab + af + ag + aj + bg + ef + ei + fi + fj + a}}{(q)_a(q)_b(q)_e(q)_f(q)_g(q)_i(q)_j(q)_a+g(q)_a+j(q)_b+g(q)_e+i(q)_f+i} \\
&\times \frac{q^{b+e+f}}{(q)_{f+j}}.
\end{aligned}$$

Now, (3.8) follows from (2.12) after letting $(a, b, e, f, g, i, j) \rightarrow (a, b, c, d, f, g, e)$.

For $\Phi_{10_5}(q)$, it suffices to prove

$$\begin{aligned}
S_{10_5}(q) &:= \sum_{a,b,c,d,e,f,g,i,j,k \geq 0} (-1)^{j+k} \frac{q^{\frac{j(3j+1)}{2} + \frac{k(7k+5)}{2} + i^2 + ab + ai + aj + bc + bj + bk + cd + ck + de + dk + ef + ek + fg}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_i(q)_j(q)_k(q)_a+i(q)_a+j(q)_b+j} \\
&\times \frac{q^{fk+gk+a+b+c+d+e+f+g}}{(q)_{b+k}(q)_{c+k}(q)_{d+k}(q)_{e+k}(q)_{f+k}(q)_{g+k}} \\
&= \frac{1}{(q)_\infty^9} h_7.
\end{aligned} \tag{3.9}$$

Apply (2.3) to the i -sum, (2.6) with $n = 3$ to the j -sum, (2.1) to the a -sum and simplify, then (2.2) to the j -sum to obtain

$$\begin{aligned}
S_{10_5}(q) &= \frac{1}{(q)_\infty^2} \sum_{b,c,d,e,f,g,k \geq 0} (-1)^k \frac{q^{\frac{k(7k+5)}{2} + bc + bk + cd + ck + de + dk + ef + ek + fg + fk + gk + b + c + d + e + f}}{(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_k(q)_{b+k}(q)_{c+k}(q)_{d+k}(q)_{e+k}(q)_{f+k}} \\
&\times \frac{q^g}{(q)_{g+k}}.
\end{aligned}$$

Now, (3.9) follows from (2.10) after letting $k \rightarrow a$.

For $\Phi_{-10_8}(q)$, it suffices to prove

$$\begin{aligned}
 S_{-10_8}(q) &:= \sum_{a,b,c,d,e,f,g,h,i,k \geq 0} (-1)^i \frac{q^{\frac{i(5i+3)}{2} + k(3k+2) + ab + ae + ai + ak + bc + bi + cd + ci + di + ef + ek + fg + fk + gh}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+i}(q)_{a+k}(q)_{b+i}} \\
 &\times \frac{q^{gk+hk+a+b+c+d+e+f+g+h}}{(q)_{c+i}(q)_{d+i}(q)_{e+k}(q)_{f+k}(q)_{g+k}(q)_{h+k}} \\
 &= \frac{1}{(q)_{\infty}^{10}} h_5 h_6.
 \end{aligned} \tag{3.10}$$

We now have

$$\begin{aligned}
 S_{-10_8}(q) &= \frac{1}{(q)_{\infty}} \sum_{a,b,c,d,e,f,g,h,i,k,j,l \geq 0} (-1)^{i+l} \frac{q^{\frac{3i(i+1)}{2} + j^2 + j + \frac{l(l+1)}{2} + 2ij + il + jl + k(3k+2) + ab + ae + ak + bc + bi}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_l} \\
 &\times \frac{q^{cd+c(i+j)+d(i+j+l)+ef+ek+fg+fk+gh+gk+hk+a+b+c+d+e+f+g+h}}{(q)_{a+i}(q)_{a+k}(q)_{e+k}(q)_{f+k}(q)_{g+k}(q)_{h+k}(q)_{b+i+j}(q)_{c+i+j+l}} \\
 &\text{(apply (2.6) to the } i\text{-sum with } n = 5\text{)} \\
 &= \frac{1}{(q)_{\infty}^4} \sum_{a,e,f,g,h,i,k,j,l \geq 0} (-1)^{i+l} \frac{q^{\frac{3i(i+1)}{2} + j^2 + j + \frac{l(l+1)}{2} + 2ij + il + jl + k(3k+2) + ae + ak + ef + ek + fg + fk + gh + hk}}{(q)_a(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_l(q)_{a+k}(q)_{e+k}(q)_{f+k}} \\
 &\times \frac{q^{a+e+f+g+h}}{(q)_{g+k}(q)_{h+k}}
 \end{aligned}$$

(evaluate the d -sum, c -sum and b -sum with (2.1) and simplify)

$$= \frac{1}{(q)_{\infty}^4} h_5 \sum_{a,e,f,g,h,k \geq 0} \frac{q^{k(3k+2) + ak + ek + fk + gk + hk + ae + ef + fg + gh + a + e + f + g + h}}{(q)_a(q)_e(q)_f(q)_g(q)_h(q)_k(q)_{a+k}(q)_{e+k}(q)_{f+k}(q)_{g+k}(q)_{h+k}}$$

(evaluate the ijl -sum using (2.5)).

Now, (3.10) follows from (1.1) after applying $(a, e, f, g, h, k) \rightarrow (c, d, e, f, g, a)$.

For $\Phi_{10_{10}}(q)$, it suffices to prove

$$\begin{aligned}
 S_{10_{10}}(q) &:= \sum_{a,c,d,e,f,g,h,i,j,k \geq 0} (-1)^{i+j} \frac{q^{\frac{i(3i+1)}{2} + \frac{j(3j+1)}{2} + k(3k+2) + ah + ai + cd + ck + de + dk + ef + ek + fg + fk + gh}}{(q)_a(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+i}(q)_{c+k}(q)_{d+k}} \\
 &\times \frac{q^{gj+gk+hi+hj+a+c+d+e+f+g+h}}{(q)_{e+k}(q)_{f+k}(q)_{g+k}(q)_{g+j}(q)_{h+j}(q)_{h+i}} \\
 &= \frac{1}{(q)_{\infty}^8} h_6.
 \end{aligned} \tag{3.11}$$

Apply (2.6) with $n = 3$ to the i -sum, (2.1) to the a -sum and simplify, (2.2) to the i and simplify to obtain

$$\begin{aligned} S_{10_{10}}(q) &= \frac{1}{(q)_\infty} \sum_{c,d,e,f,g,h,j,k \geq 0} (-1)^j \frac{q^{\frac{j(3j+1)}{2} + k(3k+2) + cd + ck + de + dk + ef + ek + fg + fk + gh + gj + gk + hj + c}}{(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_j(q)_k(q)_{c+k}(q)_{d+k}(q)_{e+k}(q)_{f+k}} \\ &\times \frac{q^{d+e+f+g+h}}{(q)_{g+k}(q)_{g+j}(q)_{h+j}}. \end{aligned}$$

Now, (3.11) follows from (2.13) after letting $(c, d, e, f, g, h, j, k) \rightarrow (h, g, f, e, d, c, b, a)$.

For $\Phi_{10_{15}}(q)$, it suffices to prove

$$\begin{aligned} S_{10_{15}}(q) &:= \sum_{a,b,c,d,e,g,h,i,j,k \geq 0} (-1)^{i+j} \frac{q^{\frac{i(5i+3)}{2} + \frac{j(5j+3)}{2} + k^2 + ab + ah + ai + bc + bi + bj + cd + cj + de + dj + ej + gh + gi}}{(q)_a(q)_b(q)_c(q)_d(q)_e(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+i}(q)_{b+i}(q)_{b+j}} \\ &\times \frac{q^{gk + hi + a + b + c + d + e + g + h}}{(q)_{c+j}(q)_{d+j}(q)_{e+j}(q)_{g+i}(q)_{g+k}(q)_{h+i}} \\ &= \frac{1}{(q)_\infty^{10}} h_5^2. \end{aligned} \tag{3.12}$$

Apply (2.3) to the k -sum, (2.6) with $n = 5$ to the j -sum, (2.1) to the e -sum and simplify, to the d -sum and simplify and to the c -sum and simplify and (2.5) to obtain

$$S_{10_{15}}(q) = \frac{1}{(q)_\infty^5} h_5 \sum_{a,b,g,h,i \geq 0} (-1)^i \frac{q^{\frac{i(5i+3)}{2} + ab + ah + ai + bi + gh + gi + hi + a + b + g + h}}{(q)_a(q)_b(q)_g(q)_h(q)_i(q)_{a+i}(q)_{b+i}(q)_{g+i}(q)_{h+i}}.$$

Now, (3.12) follows from (2.8) after letting $(a, b, g, h, i) \rightarrow (c, b, e, d, a)$.

For $\Phi_{10_{19}}(q)$, it suffices to prove

$$\begin{aligned} S_{10_{19}}(q) &:= \sum_{a,c,d,e,f,g,h,i,j,k \geq 0} (-1)^{j+k} \frac{q^{i(2i+1) + \frac{j(3j+1)}{2} + \frac{k(5k+3)}{2} + ah + ai + cd + ck + de + dek + ef + ek + fg + fk}}{(q)_a(q)_c(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+i}(q)_{c+k}(q)_{d+k}} \\ &\times \frac{q^{fj + gh + gi + gj + hi + a + c + d + e + f + g + h}}{(q)_{e+k}(q)_{f+k}(q)_{f+j}(q)_{g+j}(q)_{g+i}(q)_{h+i}} \\ &= \frac{1}{(q)_\infty^9} h_4 h_5. \end{aligned} \tag{3.13}$$

Apply (2.6) with $n = 5$ to the k -sum, (2.1) to the c -sum and simplify, to the d -sum and simplify and to the e -sum and simplify and (2.5) to obtain

$$S_{10_{19}}(q) = \frac{1}{(q)_\infty^4} \sum_{a,f,g,h,i,j \geq 0} (-1)^j \frac{q^{i(2i+1) + \frac{j(3j+1)}{2} + ah + ai + fg + fj + gh + gi + gj + hi + a + f + g + h}}{(q)_a(q)_f(q)_g(q)_h(q)_i(q)_j(q)_{a+i}(q)_{f+j}(q)_{g+j}(q)_{g+i}(q)_{h+i}}.$$

Now, (3.13) follows from (2.9) after letting $(a, f, g, h, i, j) \rightarrow (a, d, c, b, f, e)$.

For $\Phi_{10_{26}}(q)$, it suffices to prove

$$\begin{aligned}
 S_{10_{26}}(q) &:= \sum_{a,b,c,e,f,g,h,i,j,k \geq 0} (-1)^i \frac{q^{h(2h+1)+\frac{i(3i+1)}{2}+j^2+k(2k+1)+ab+ag+ah+ai+bc+bh+ch+ef+ek+fg}}{(q)_a(q)_b(q)_c(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+h}(q)_{a+i}(q)_{b+h}} \\
 &\times \frac{q^{fk+gi+gj+gk+a+b+c+e+f+g}}{(q)_{c+h}(q)_{e+k}(q)_{f+k}(q)_{g+i}(q)_{g+j}(q)_{g+k}} \\
 &= \frac{1}{(q)_\infty^9} h_4^2.
 \end{aligned} \tag{3.14}$$

Apply (2.3) to the j -sum, (2.6) with $n = 4$ to the k -sum, (2.1) to the e -sum and simplify and to the f -sum and simplify and (2.4) to obtain

$$S_{10_{26}}(q) = \frac{1}{(q)_\infty^4} h_4 \sum_{a,b,c,g,h,i \geq 0} (-1)^i \frac{q^{h(2h+1)+\frac{i(3i+1)}{2}+ab+ag+ah+ai+bc+bh+ch+gi+a+b+c+g}}{(q)_a(q)_b(q)_c(q)_g(q)_h(q)_i(q)_{a+h}(q)_{a+i}(q)_{b+h}(q)_{c+h}(q)_{g+i}}.$$

Now, (3.14) follows from (2.9) after letting $(a, b, c, g, h, i) \rightarrow (c, b, a, d, f, e)$.

For $\Phi_{10_{28}}(q)$, it suffices to prove

$$\begin{aligned}
 S_{10_{28}}(q) &:= \sum_{a,b,d,e,f,g,h,i,j,k \geq 0} (-1)^{i+j} \frac{q^{\frac{i(3i+1)}{2}+\frac{j(5j+3)}{2}+k(2k+1)+ab+ah+ai+aj+bi+de+dk+ef+ek+fg+ff}}{(q)_a(q)_b(q)_d(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+i}(q)_{a+j}(q)_{b+i}} \\
 &\times \frac{q^{fk+gh+gj+hj+a+b+d+e+f+g+h}}{(q)_{d+k}(q)_{e+k}(q)_{f+j}(q)_{f+k}(q)_{g+j}(q)_{h+j}} \\
 &= \frac{1}{(q)_\infty^9} h_4 h_5.
 \end{aligned} \tag{3.15}$$

Apply (2.6) with $n = 3$ to the i -sum, (2.1) to the b -sum and simplify and (2.2) to the i -sum to obtain

$$\begin{aligned}
 S_{10_{28}}(q) &= \frac{1}{(q)_\infty} \sum_{a,d,e,f,g,h,j,k \geq 0} (-1)^j \frac{q^{\frac{j(5j+3)}{2}+k(2k+1)+ah+aj+de+dk+ef+ek+fg+fj+fk+gh+gj+hj}}{(q)_a(q)_d(q)_e(q)_f(q)_g(q)_h(q)_j(q)_k(q)_{a+j}(q)_{d+k}(q)_{e+k}(q)_{f+j}} \\
 &\times \frac{q^{a+d+e+f+g+h}}{(q)_{f+k}(q)_{g+j}(q)_{h+j}}.
 \end{aligned}$$

Now, (3.15) follows from (2.14) after letting $(a, d, e, f, g, h, j, k) \rightarrow (f, a, b, c, d, e, g, h)$.

For $\Phi_{10_{44}}(q)$, it suffices to prove

$$\begin{aligned}
S_{1044}(q) &:= \sum_{a,b,c,e,f,g,h,i,j,k \geq 0} (-1)^{h+j+k} \frac{q^{\frac{h(3h+1)}{2} + i(2i+1) + \frac{j(3j+1)}{2} + \frac{k(3k+1)}{2} + ab + ag + ai + aj + bc + bj + bk + ck}}{(q)_a(q)_b(q)_c(q)_e(q)_f(q)_g(q)_h(q)_i(q)_j(q)_k(q)_{a+i}(q)_{a+j}(q)_{b+j}} \\
&\times \frac{q^{ef+eh+fg+fh+fi+gi+a+b+c+e+f+g}}{(q)_{b+k}(q)_{c+k}(q)_{e+h}(q)_{f+h}(q)_{f+i}(q)_{g+i}} \\
&= \frac{1}{(q)_\infty^7} h_4.
\end{aligned} \tag{3.16}$$

Apply (2.6) with $n = 3$ to the h -sum, (2.1) to the e -sum and simplify, (2.2) to the h -sum, (2.6) with $n = 3$ to the k -sum, (2.1) to the c -sum and simplify and (2.2) to the k -sum to obtain

$$S_{1044}(q) = \frac{1}{(q)_\infty^2} \sum_{a,b,f,g,i,j \geq 0} (-1)^j \frac{q^{i(2i+1) + \frac{j(3j+1)}{2} + ab + ag + ai + aj + bj + fg + fi + gi + a + b + f + g}}{(q)_a(q)_b(q)_f(q)_g(q)_i(q)_j(q)_{a+i}(q)_{a+j}(q)_{b+j}(q)_{f+i}(q)_{g+i}}.$$

Now, (3.16) follows from (2.9) after letting $(a, b, f, g, i, j) \rightarrow (c, d, a, b, f, e)$. \square

ACKNOWLEDGEMENTS

The second author would like to thank Don Zagier for his question posed during the workshop “Low-dimensional topology and number theory”, August 17–23, 2014 at Oberwolfach and the organizers (in particular, Frits Beukers) of the conference “Automatic sequences, Number Theory, Aperiodic Order”, October 28–30, 2015 at TU Delft for the opportunity to discuss these results. He also thanks Jeremy Lovejoy for his helpful comments and suggestions.

REFERENCES

- [1] C. Armond, *The head and tail conjecture for alternating knots*, Algebr. Geom. Topol. **13** (2013), no. 5, 2809–2826.
- [2] C. Armond, O. Dasbach, *Rogers-Ramanujan type identities and the head and tail of the colored Jones polynomial*, preprint available at <http://arxiv.org/abs/1106.3948>
- [3] O. Dasbach, X.-S. Lin, *On the head and the tail of the colored Jones polynomial*, Compos. Math. **142** (2006), no. 5, 1332–1342.
- [4] M. Elhamdadi, M. Hajij, *Pretzel knots and q-series*, preprint available at <http://arxiv.org/abs/1512.00129>
- [5] S. Garoufalidis, T. Lê, *Nahm sums, stability and the colored Jones polynomial*, Res. Math. Sci. **2** (2015), Art. 1, 55pp.
- [6] S. Gukov, S. Nawata, I. Saberi, M. Stosic and Piotr Sulkowski, *Sequencing BPS spectra*, preprint available at <http://arxiv.org/abs/1512.07883>
- [7] M. Hajij, *The colored Kauffman skein relation and the head and tail of the colored Jones polynomial*, preprint available at <http://arxiv.org/abs/1401.4537>
- [8] A. Keilthy, R. Osburn, *Rogers-Ramanujan type identities for alternating knots*, J. Number Theory, **161** (2016), 255–280.
- [9] S. Nawata, A. Oblomkov, *Lectures on knot homology*, preprint available at <http://arxiv.org/abs/1510.01795>
- [10] L. Rozansky, *Khovanov homology of a unicolored B-adequate link has a tail*, Quantum Topol. **5** (2014), no. 4, 541–579.

- [11] R. van der Veen, *The degree of the colored HOMFLY polynomial*, preprint available at <http://arxiv.org/abs/1501.00123>

72 OAKTREE ROAD, MERVILLE, STILLORGAN, COUNTY DUBLIN

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY COLLEGE DUBLIN, BELFIELD, DUBLIN 4, IRELAND

E-mail address: paul.beirne@ucdconnect.ie

E-mail address: robert.osburn@ucd.ie