

RESURGENCE, HABIRO ELEMENTS AND STRANGE IDENTITIES

SAMUEL CREW, VERONICA FANTINI, ANKUSH GOSWAMI, ROBERT OSBURN,
AND CAMPBELL WHEELER

ABSTRACT. We prove resurgence properties for the Borel transform of a formal power series associated to elements in the Habiro ring that come from radial limits of partial theta series via strange identities. As an application, we prove a conjecture in quantum topology due to Costin and Garoufalidis for two families of torus knots.

1. INTRODUCTION

The modern theory of resurgence began with the voluminous works of Écalle in 1981 [16, 17] and 1985 [18]. This theory now plays a vital rôle in a remarkable number of diverse areas, to name just a few examples: nonlinear systems of ODEs and difference equations [11, 42], algebraic combinatorics [9, 10], enumerative combinatorics and quantum field theory [8, 46], period integrals and string theory scattering amplitudes [15], wall-crossing phenomena [43] and matrix models [48]. In this paper, we are interested in a conjecture due to Costin and Garoufalidis [13] which connects resurgence and quantum topology. For other such interactions, see [3, 19, 24–26] and the references therein.

Let us recall the general setup from [13]. For further details, see [12, 14, 47, 49] or [52, Section 2] for an excellent concise review of the basic notions of resurgence and alien calculus. A formal power series

$$\mathcal{F}(x) = \sum_{n=0}^{\infty} a_n x^{-n} \in \mathbb{C}[[1/x]] \quad (1.1)$$

is called Gevrey-1 if there exists positive real numbers A, B such that

$$|a_n| \leq AB^n n!$$

for all $n \geq 0$. Consider a Gevrey-1 formal power series given by (1.1) and its Borel transform $\mathcal{B} : \mathbb{C}[[1/x]] \rightarrow \mathbb{C}[[p]]$ defined by

$$\mathcal{B}\left(\sum_{n=0}^{\infty} a_n x^{-n}\right) = a_0 \delta + \sum_{n=1}^{\infty} a_n \frac{p^{n-1}}{(n-1)!} =: a_0 \delta + G(p) \quad (1.2)$$

where δ is a formal symbol that has Laplace transform given by the constant function 1. For simplicity, we write $\mathcal{B}[\mathcal{F}](p)$ for the transform (1.2). Here, $G(p)$ has a positive radius of convergence as it arose from a Gevrey-1 power series. In a standard abuse of notation, $G(p)$ denotes the formal power series in (1.2) (and is also commonly referred to in the literature as the Borel transform of $\mathcal{F}(x)$) and the analytic continuation of the associated germ at the origin. In addition,

Date: November 29, 2024.

2020 Mathematics Subject Classification. 57K16, 11F37, 30B40, 40A05.

Key words and phrases. Resurgence, Habiro ring, quantum topology, strange identities.

$G(p)$ is called a *resurgent function* if it is endlessly analytically continuable [49]. This property means $G(p)$ extends to a (possibly multi-valued) holomorphic function along unbounded paths that need only circumvent a discrete set of singularities. In the present work, we consider the case where $G(p)$ defines a multi-valued function on $\mathbb{C} \setminus \mathcal{N}$, with \mathcal{N} a discrete set of singularities lying on the ray $\mathbb{R}_{>0}$, e.g., $\mathcal{N} = \left\{ \frac{n^2\pi^2}{6} : n \in \mathbb{Z}_{>0} \right\}$.

Assuming $G(p)$ satisfies suitable growth conditions, the left and right Borel resummations of $\mathcal{F}(x)$ are defined as

$$S^+[\mathcal{F}](x) = \int_{\gamma_l} e^{-px} \mathcal{B}[\mathcal{F}](p) dp$$

and

$$S^-[\mathcal{F}](x) = \int_{\gamma_r} e^{-px} \mathcal{B}[\mathcal{F}](p) dp$$

where $x \in \mathbb{C}$ and γ_l and γ_r are contours in $\mathbb{C} \setminus \mathcal{N}$ from 0 to ∞ that turn left (respectively, right) at each singularity in \mathcal{N} (see Figure 1). In addition, the left and right Borel resummations give analytic functions on an sector of opening angle π in the complex x -plane.




FIGURE 1. A singularity set $\mathcal{N} \subset \mathbb{C}$ and the contours γ_l and γ_r .

Following [13, Sections 4.2–4.3], the median resummation of \mathcal{F} is given by

$$S^{\text{med}}(x) = S^{\text{med}}[\mathcal{F}](x) := \frac{1}{2} \left(S^+[\mathcal{F}](x) + S^-[\mathcal{F}](x) \right). \quad (1.3)$$

In general, care is required when defining median summations, see [32, Section 2]. Under favorable circumstances, for example when $G(p)$ is the Borel transform of a formal (Gevrey-1) solution in powers of $1/x$ to a linear differential equation, each of $S^\pm[\mathcal{F}](x)$ (and thus, of course, $S^{\text{med}}(x)$) is a well-defined homomorphic solution. This is one incarnation of the terminology “resummation”.

In this paper, we will study the median resummation of the Borel transform for certain formal power series associated to knots.

Let K be a knot and $J_N(K; q)$ be the usual colored Jones polynomial, normalized to be 1 for the unknot. If $N = 2$, then we recover the Jones polynomial [39]. As a knot invariant, the colored Jones polynomial is of fundamental importance in several open problems in quantum topology (see, e.g., [20, 22, 40, 51, 55]). We now recall the Habiro ring [31]

$$\mathcal{H} := \varprojlim_n \mathbb{Z}[q] / \langle (q)_n \rangle$$

where

$$(a)_n = (a; q)_n := \prod_{k=1}^n (1 - aq^{k-1})$$

is the standard q -Pochhammer symbol. Every element of \mathcal{H} may be written as

$$\sum_{n=0}^{\infty} b_n(q)(q)_n$$

where $b_n(q) \in \mathbb{Z}[q]$. Given K , there exists an element $\Phi_K(q) \in \mathcal{H}$ called the Kashaev invariant [41] which matches $J_N(K; q)$ (up to a prefactor) when $q = \zeta_N := e^{\frac{2\pi i}{N}}$ [38, Theorem 2]. To illustrate, for the trefoil knot $K = 3_1$, we have [30, 44]

$$J_N(3_1; q) = q^{1-N} \sum_{n=0}^{\infty} q^{-nN} (q^{1-N})_n. \quad (1.4)$$

In this case, the Kontsevich-Zagier series [55]

$$\Phi_{3_1}(q) := \sum_{n=0}^{\infty} (q)_n \in \mathcal{H} \quad (1.5)$$

gives the Kashaev invariant for the trefoil [41] and so from (1.4) and (1.5), we observe

$$\Phi_{3_1}(\zeta_N)\zeta_N = J_N(3_1; \zeta_N).$$

Now, a crucial aspect of the computations in [13] is the “strange identity”

$$\Phi_{3_1}(q) \text{ “} = \text{”} - \frac{1}{2} \sum_{n=1}^{\infty} n \left(\frac{12}{n} \right) q^{\frac{n^2-1}{24}} \quad (1.6)$$

where “=” means that both sides agree to infinite order at any root of unity and $\left(\frac{12}{*}\right)$ is the quadratic character of conductor 12. More precisely, replacing q by the product of a root of unity and e^{-x} , then letting $x \rightarrow 0^+$, the right-hand side has an asymptotic expansion as a power series in x and this power series is given by the left-hand side at q . For further details, see [2, 54]. If $q = e^{2\pi i x}$, $x \in \mathbb{H}$, then (1.6) implies for $0 \neq \alpha \in \mathbb{Q}$

$$e^{\frac{\pi i \alpha}{12}} \Phi_{3_1}(e^{2\pi i \alpha}) = -\frac{1}{2} \sum_{n=1}^{\infty} n \left(\frac{12}{n} \right) e^{\frac{n^2 \pi i \alpha}{12}} \quad (1.7)$$

where the right-hand side of (1.7) is interpreted as the radial limit $x \rightarrow \alpha$. The significance of “identities” akin to (1.6) is evident in various applications, e.g., asymptotics and congruences for Fishburn and generalized Fishburn numbers modulo prime powers [1, 2, 4, 6, 27, 28, 53, 54], the quantum modularity of $\Phi_{3_1}(q)$ [29, 55] and expressing WRT invariants of Brieskorn homology spheres in terms of limiting values of Eichler integrals [33–35]. For a recent advance using the “Bailey machinery” which not only recovers (1.6), but produces a wealth of new examples, see [45].

Elements of \mathcal{H} can also be formally expanded around any root of unity. Therefore, we let

$$F_K(x) := \Phi_K(e^{-\frac{1}{x}}) \in \mathbb{C}[[1/x]]. \quad (1.8)$$

To emphasize the dependence on K , we write $G_K(p)$ for $G(p)$ and $S_K^{\text{med}}(x)$ for $S^{\text{med}}(x)$. Finally, we use the notation \doteq to denote equality up to a prefactor which depends on K . We can now state the main conjecture from [13] (slightly edited for clarity) which is part of a larger program to understand the analytic continuation of invariants of “knotted objects” arising in Chern-Simons theory [21, 23].

Conjecture 1.1. *For every knot K ,*

- (1) $F_K(x)$ has a resurgent Borel transform $G_K(p)$.
- (2) $S_K^{\text{med}}(x)$ is an analytic function defined on $\Re(x) > 0$ with radial limits at the points $\frac{1}{2\pi i}\mathbb{Q}$ of its natural boundary.
- (3) If (2) is true, then consider the radial limit

$$S_K^{\text{med}}\left(-\frac{1}{2\pi i\alpha}\right) := \lim_{x \rightarrow -\frac{1}{2\pi i\alpha}} S_K^{\text{med}}(x)$$

where $\Re(x) > 0$. For $0 \neq \alpha \in \mathbb{Q}$, we have $S_K^{\text{med}}\left(-\frac{1}{2\pi i\alpha}\right) \doteq \Phi_K(e^{2\pi i\alpha})$.

In [13, Theorem 2.5, Theorem 3.1], Conjecture 1.1 (1) and (2) were proven for $K = 3_1$. Here,

$$F_{3_1}(x) = e^{-\frac{1}{24x}} \sum_{n=0}^{\infty} (e^{-\frac{1}{x}})_n$$

where an adjustment of (1.8) has been made. The Borel transform $G_{3_1}(p)$ of $F_{3_1}(x)$ is explicitly given by [13, Theorem 3.1]

$$G_{3_1}(p) = \frac{3\pi}{2\sqrt{2}} \sum_{n=1}^{\infty} \frac{n \binom{12}{n}}{\left(-p + \frac{n^2\pi^2}{6}\right)^{5/2}}. \quad (1.9)$$

Moreover, a close inspection of the proofs of Theorems 2.5 and 3.1 in [13] reveals that it is actually (1.6) and the periodic function $\left(\frac{12}{*}\right)$ which determine the analytic nature of $G_{3_1}(p)$ and $S_{3_1}^{\text{med}}(x)$. Our main goal in this paper is to prove resurgence properties for formal power series associated to elements in \mathcal{H} that satisfy a general type of strange identity motivated by (1.6). Here, we emphasize the importance of periodic functions. Before stating our main result, we introduce some notation.

Let $k_1, k_2 \in \mathbb{Z}$ with $k_1 < k_2$, $M \geq 2$ be an integer, $0 \neq c \in \mathbb{R}$ and f be the function

$$f(n) := \begin{cases} c & \text{if } n \equiv k_1, M - k_1 \pmod{M}, \\ -c & \text{if } n \equiv k_2, M - k_2 \pmod{M}, \\ 0 & \text{otherwise.} \end{cases} \quad (1.10)$$

Note that f is an even function of period M with mean value zero. For integers $a \geq 0$ and $b > 0$, consider the partial theta series

$$\theta_{a,b,f}^{(\nu)}(q) := \sum_{n=0}^{\infty} n^\nu f(n) q^{\frac{n^2-a}{b}} \quad (1.11)$$

where $q = e^{2\pi i x}$, $x \in \mathbb{H}$, and $\nu \in \{0, 1\}$. For $\ell \in \mathbb{Z}$, set

$$\tilde{f}(\ell) := (-1)^\ell \sin\left(\frac{(k_2 - k_1)\ell\pi}{M}\right) \sin\left(\frac{(M - k_1 - k_2)\ell\pi}{M}\right). \quad (1.12)$$

Suppose there exists

$$\Phi_f(q) := \sum_{n=0}^{\infty} A_{n,f}(q)(q)_n \in \mathcal{H}$$

where $A_{n,f}(q) \in \mathbb{Z}[q]$ such that

$$\Phi_f(q) \text{ “} = \text{”} \theta_{a,b,f}^{(1)}(q). \quad (1.13)$$

Finally, define

$$\mathcal{F}_f(x) := e^{-\frac{a}{bx}} \Phi_f(e^{-\frac{1}{x}}) = \sum_{n=0}^{\infty} \frac{C_{n,f}}{n!} \left(\frac{1}{bx}\right)^n. \quad (1.14)$$

The asymptotics of the partial theta series $\theta_{a,b,f}^{(1)}(q)$ around $q = 1$, which by the strange identity in (1.13) gives the expansion of $\Phi_f(q)$, can be computed by L -values. By standard calculations using the Mellin transform (see, e.g., [2, 28, 55]), we find that

$$C_{n,f} = (-1)^n L(-2n - 1, f)$$

where

$$L(s, f) := \sum_{n=1}^{\infty} \frac{f(n)}{n^s}$$

the L -series associated to f . In particular, using [2, Lemma 3.2], we obtain that

$$C_{n,f} = (-1)^{n+1} \frac{M^{2n+1}}{2n+2} \sum_{m=1}^M f(m) B_{2n+2} \left(\frac{m}{M}\right) \quad (1.15)$$

where for $k \geq 0$, $B_k(x)$ denotes the k th Bernoulli polynomial. By (1.14) and (1.15), we have

$$\mathcal{F}_f \left(\frac{1}{x}\right) = \sum_{n=0}^{\infty} \frac{(-1)^{n+1} M^{2n+1}}{(2n+2)n!} \sum_{m=1}^M f(m) B_{2n+2} \left(\frac{m}{M}\right) \left(\frac{x}{b}\right)^n. \quad (1.16)$$

We will denote the Borel transform of $\mathcal{F}_f(x)$ by $G_f(p)$. Here, we write $S_f^{\text{med}}(x)$ for $S^{\text{med}}(x)$,

$$S_f^{\text{med}} \left(-\frac{1}{2\pi i \alpha}\right) := \lim_{x \rightarrow -\frac{1}{2\pi i \alpha}} S_f^{\text{med}}(x),$$

$\theta_{a,b,f}^{(\nu)}(x)$ for $\theta_{a,b,f}^{(\nu)}(q)$ and $\theta_{a,b,f}^{(\nu)}(\alpha) := \lim_{x \rightarrow \alpha} \theta_{a,b,f}^{(\nu)}(x)$ where $0 \neq \alpha \in \mathbb{Q}$. Our main result is now the following.

Theorem 1.2. *For f given by (1.10) and $\mathcal{F}_f(x)$ as in (1.16), we have the following:*

- (1) $\mathcal{F}_f(x)$ has a resurgent Borel transform $G_f(p)$.
- (2) $S_f^{\text{med}}(x)$ is an analytic function defined on $\Re(x) > 0$ with radial limits at the points $\frac{1}{2\pi i} \mathbb{Q}$ of its natural boundary.
- (3) For $0 \neq \alpha \in \mathbb{Q}$, we have

$$S_f^{\text{med}} \left(-\frac{1}{2\pi i \alpha}\right) = \frac{cbe^{\pi i/4}}{M\pi(i\alpha)^{3/2}} \int_0^{+\infty} \frac{\theta_{0,4M^2,\tilde{f}}^{(0)}(bp)}{\left(\frac{1}{\alpha} + p\right)^{3/2}} dp + \left(\frac{b}{i\alpha}\right)^{3/2} \frac{\sqrt{2}c}{M^2} \theta_{0,4M^2,\tilde{f}}^{(1)} \left(-\frac{b}{\alpha}\right). \quad (1.17)$$

Remark 1.3. The proof of Theorem 1.2 demonstrates resurgence properties for the normalized partial theta series $e^{-\frac{a}{bx}}\theta_{a,b,f}^{(1)}(e^{-\frac{1}{x}})$ without assuming (1.13). Our results and those in the announcement [32] appear to be related; however, we give a proof that can be used to prove Conjecture 1.1 for our cases of interest (see Corollary 1.5).

We give two applications of Theorem 1.2. The first result explicitly computes (1.17) in the following situation. For coprime positive integers s and t , $1 \leq m \leq t-1$ and $1 \leq n \leq s-1$, let $\chi_{2st}^{(n,m)}(k)$ be the periodic function obtained by choosing $c = 1$, $M = 2st$, $k_1 = nt - ms$ and $k_2 = nt + ms$ in (1.10), i.e.,

$$\chi_{2st}^{(n,m)}(k) = \begin{cases} 1 & \text{if } k \equiv \pm(nt - ms) \pmod{2st}, \\ -1 & \text{if } k \equiv \pm(nt + ms) \pmod{2st}, \\ 0 & \text{otherwise.} \end{cases} \quad (1.18)$$

For coprime odd integers s, t , define the sets

$$\mathcal{D}_1(s, t) := \left\{ (n, m) : 1 \leq n \leq \frac{s-1}{2}, 1 \leq m \leq t-1 \right\} \quad (1.19)$$

and

$$\mathcal{D}_2(s, t) := \left\{ (n, m) : 1 \leq n \leq s-1, 1 \leq m \leq \frac{t-1}{2} \right\}. \quad (1.20)$$

There is a bijection between $\mathcal{D}_1(s, t)$ and $\mathcal{D}_2(s, t)$ (see Lemma 3.1 and (3.2)). For s and t coprime and of opposite parities, we consider

$$\mathcal{D}_3(s, t) := \begin{cases} \{(n, m) : 1 \leq n \leq \frac{s-1}{2}, 1 \leq m \leq t-1\} & \text{if } s \equiv 1 \pmod{2}, s \not\equiv t \pmod{2}, \\ \{(n, m) : 1 \leq n \leq s-1, 1 \leq m \leq \frac{t-1}{2}\} & \text{if } s \equiv 0 \pmod{2}, s \not\equiv t \pmod{2}. \end{cases}$$

Finally, define the set $\mathcal{D}(s, t)$ of distinct pairs as follows:

$$\mathcal{D}(s, t) := \begin{cases} \mathcal{D}_1(s, t) \text{ or } \mathcal{D}_2(s, t) & \text{if } s \equiv t \equiv 1 \pmod{2}, \\ \mathcal{D}_3(s, t) & \text{if } s \not\equiv t \pmod{2}. \end{cases} \quad (1.21)$$

One sees that $D(s, t) := |\mathcal{D}(s, t)| = \frac{(s-1)(t-1)}{2}$.

Theorem 1.4. *Let s and t be coprime positive integers and $(n, m) \in \mathcal{D}(s, t)$. For $0 \neq \alpha \in \mathbb{Q}$, we have*

$$S_{\chi_{2st}^{(n,m)}}^{\text{med}} \left(-\frac{1}{2\pi i \alpha} \right) = \theta_{0,4st, \chi_{2st}^{(n,m)}}^{(1)}(\alpha). \quad (1.22)$$

The second result verifies Conjecture 1.1 for two families of torus knots.

Corollary 1.5. *Let $u, k \in \mathbb{Z}_{\geq 1}$. Conjecture 1.1 is true for the families of torus knots $T(2, 2u+1)$ and $T(3, 2^k)$.*

Corollary 1.5 generalizes Theorems 2.5 and 3.1 in [13] and upon taking either $u = 1$ or $k = 1$ confirms Conjecture 1.1 for the trefoil knot $3_1 = T(2, 3) = T(3, 2)$.

The paper is organized as follows. In Section 2, we prove Theorem 1.2 by first providing an explicit evaluation of $G_f(p)$ in (2.9), which generalizes (1.9), and then computing $S_f^{\text{med}}(x)$ directly via a careful contour deformation argument. In Section 3, we prove Theorem 1.4 and Corollary 1.5. In Section 4, we make some concluding remarks.

2. PROOF OF THEOREM 1.2

Proof of Theorem 1.2. The Borel transform of $\mathcal{F}_f(1/x)$ from (1.16) is given by

$$\mathcal{B}[\mathcal{F}_f](p) = C_M \delta + G_f(p)$$

where

$$C_M = -\frac{M}{2} \sum_{m=1}^M f(m) B_2 \left(\frac{m}{M} \right)$$

and

$$\begin{aligned} G_f(p) &= \sum_{n=1}^{\infty} \frac{(-1)^{n+1} M^{2n+1}}{(2n+2)n!} \sum_{m=1}^M f(m) B_{2n+2} \left(\frac{m}{M} \right) \frac{p^{n-1}}{(n-1)! b^n} \\ &= M \sum_{n=0}^{\infty} \sum_{m=1}^M \frac{(-1)^n f(m) B_{2n+4} \left(\frac{m}{M} \right)}{(2n+4)!} \frac{(2n+3)!}{n!(n+1)!} \left(\frac{M^2}{b} \right)^{n+1} p^n. \end{aligned} \quad (2.1)$$

From the generating function

$$\frac{x e^{xt}}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n(t)}{n!} x^n,$$

where $|x| < 2\pi$, we deduce

$$\sum_{n=0}^{\infty} \frac{B_{2n}(t)}{(2n)!} x^{2n} = \frac{x}{2} \left(\frac{e^{xt}}{e^x - 1} - \frac{e^{-xt}}{e^{-x} - 1} \right) = \frac{x(e^{xt} + e^{-(t-1)x})}{2(e^x - 1)}. \quad (2.2)$$

Assuming $|y| < \frac{2\pi}{M}$, $x = iMy$, and taking $t = \frac{m}{M}$, where $1 \leq m \leq M$, we use (2.2) to obtain

$$\begin{aligned} \frac{1}{My} \sum_{m=1}^M f(m) \sum_{n=0}^{\infty} \frac{B_{2n+2} \left(\frac{m}{M} \right)}{(2n+2)!} (iMy)^{2n+2} &= \frac{i}{2(e^{iMy} - 1)} \sum_{m=1}^M f(m) \left(e^{imy} + e^{-i(m-M)y} \right) \\ &= -2c \cdot \frac{\sin \left(\frac{(k_2 - k_1)y}{2} \right) \sin \left(\frac{(M - k_1 - k_2)y}{2} \right)}{\sin \left(\frac{My}{2} \right)}. \end{aligned} \quad (2.3)$$

If we let $y = \sqrt{p}$ in (2.3), then

$$\begin{aligned} g_{1,f}(p) &:= M^3 \sum_{n=0}^{\infty} \sum_{m=1}^M f(m) \frac{B_{2n+4} \left(\frac{m}{M} \right)}{(2n+4)!} (-M^2 p)^n \\ &= \frac{1}{Mp^2} \sum_{n=0}^{\infty} \sum_{m=1}^M f(m) \frac{B_{2n+2} \left(\frac{m}{M} \right)}{(2n+2)!} (-M^2 p)^{n+1} - \frac{C_M}{p} \\ &= -\frac{1}{p^{3/2}} \left(2c \cdot \frac{\sin \left(\frac{(k_2 - k_1)\sqrt{p}}{2} \right) \sin \left(\frac{(M - k_1 - k_2)\sqrt{p}}{2} \right)}{\sin \left(\frac{M\sqrt{p}}{2} \right)} + C_M \sqrt{p} \right). \end{aligned} \quad (2.4)$$

We conclude

$$G_f(p) = (g_{1,f} \circledast g_{2,f})(p)$$

where $g_{1,f}$ was computed in (2.4) and

$$g_{2,f}(p) := \sum_{n=0}^{\infty} \frac{(2n+3)!}{n!(n+1)!} \left(\frac{1}{b}\right)^{n+1} p^n = \frac{1}{b} \frac{6}{\left(1 - 4\frac{p}{b}\right)^{5/2}}$$

and \otimes denotes the Hadamard product of two formal power series

$$\left(\sum_{n=0}^{\infty} a_n p^n\right) \otimes \left(\sum_{n=0}^{\infty} b_n p^n\right) := \sum_{n=0}^{\infty} a_n b_n p^n.$$

It now follows that

$$G_f(p) = \frac{1}{2\pi i} \int_{-\gamma} g_{1,f}(s) g_{2,f}\left(\frac{p}{s}\right) \frac{ds}{s} \quad (2.5)$$

where γ is a circle with center at the origin and a small radius oriented clockwise (see Figure 2). Now, we consider the poles of $g_{1,f}(s)$. Note that there is no pole at $s = 0$. By (2.4), they are supported where $p = \frac{4\ell^2\pi^2}{M^2}$, $\ell \in \mathbb{Z}$. However,

$$\sin\left(\frac{(k_2 - k_1)\sqrt{p}}{2}\right) \sin\left(\frac{(M - k_1 - k_2)\sqrt{p}}{2}\right) = 0$$

when $p = \frac{4k^2\pi^2}{(k_2 - k_1)^2}$ or $p = \frac{4k^2\pi^2}{(M - k_2 - k_1)^2}$ for $k \in \mathbb{Z}$. Thus, if for some $k, \ell \in \mathbb{Z}$

$$\ell = \frac{Mk}{k_2 - k_1} \quad \text{or} \quad \ell = \frac{Mk}{M - k_2 - k_1}$$

which is true if and only if

$$\ell = \frac{\frac{M}{\gcd(M, k_2 - k_1)} k}{\frac{(k_2 - k_1)}{\gcd(M, k_2 - k_1)}} \quad \text{or} \quad \ell = \frac{\frac{M}{\gcd(M, M - k_2 - k_1)} k}{\frac{(M - k_2 - k_1)}{\gcd(M, M - k_2 - k_1)}},$$

then $g_{1,f}(s)$ does not have poles at $p = \frac{4\ell^2\pi^2}{M^2}$. Thus, the set of poles of $g_{1,f}(s)$ is given by

$$\mathcal{N}_{1,f} = \left\{ \frac{4\ell^2\pi^2}{M^2} : \ell \in \mathbb{N}, \frac{M}{\gcd(M, k_2 - k_1)} \nmid \ell \text{ or } \frac{M}{\gcd(M, M - k_2 - k_1)} \nmid \ell \right\}. \quad (2.6)$$

We next carefully calculate the residues of the poles of the integrand (2.5) which are given by (2.6). Take a circle γ_R with center at the origin and of large radius R which encloses a finite number of points N in $\mathcal{N}_{1,f}$. Thus,

$$\begin{aligned} \frac{1}{2\pi i} \int_{\gamma_R \cup \gamma} g_{1,f}(s) g_{2,f}\left(\frac{p}{s}\right) \frac{ds}{s} &= \sum_{\substack{p_i \in \mathcal{N}_{1,f} \\ 1 \leq i \leq N}} \text{Res}_{s \rightarrow p_i} \left(\frac{g_{1,f}(s) g_{2,f}\left(\frac{p}{s}\right)}{s} \right) \\ &= \sum_{\ell \in \mathbb{N}}^N \frac{g_{2,f}\left(\frac{pM^2}{4\ell^2\pi^2}\right)}{\left(\frac{4\ell^2\pi^2}{M^2}\right)} \end{aligned}$$

$$\begin{aligned}
& \times \lim_{s \rightarrow \frac{4\ell^2\pi^2}{M^2}} \left(s - \frac{4\ell^2\pi^2}{M^2} \right) \left[-\frac{1}{s^{3/2}} \left(2c \cdot \frac{\sin\left(\frac{(k_2-k_1)\sqrt{s}}{2}\right) \sin\left(\frac{(M-k_1-k_2)\sqrt{s}}{2}\right)}{\sin\left(\frac{M\sqrt{s}}{2}\right)} + C_M\sqrt{s} \right) \right] \\
& = -\frac{3\pi c}{bM^2} \sum_{\ell \in \mathbb{N}}^N \frac{\ell \tilde{f}(\ell)}{\left(\frac{\ell^2\pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} \tag{2.7}
\end{aligned}$$

where \sum^N means that ℓ runs over the first N elements $\frac{4\ell^2\pi^2}{M^2}$ of $\mathcal{N}_{1,f}$ and $\tilde{f}(\ell)$ is given by (1.12). It follows that

$$\left| \int_{\gamma_R} g_{1,f}(s) g_{2,f}\left(\frac{p}{s}\right) \frac{ds}{s} \right| \rightarrow 0 \tag{2.8}$$

as $R \rightarrow \infty$. Thus, (2.5), (2.7), (2.8) and the contour deformation given in Figure 2 imply that

$$G_f(p) = \frac{3\pi c}{M^2 b} \sum_{\ell=1}^{\infty} \frac{\ell \tilde{f}(\ell)}{\left(\frac{\ell^2\pi^2}{M^2} - \frac{p}{b}\right)^{5/2}}. \tag{2.9}$$




FIGURE 2. Contour deformation of $\gamma_R - \gamma$

The set of poles of $G_f(p)$ is then

$$\mathcal{N} := \left\{ \frac{b\ell^2\pi^2}{M^2} : \ell \in \mathbb{N}, \frac{M}{\gcd(M, k_2 - k_1)} \nmid \ell \text{ or } \frac{M}{\gcd(M, M - k_2 - k_1)} \nmid \ell \right\}.$$

Since $G_f(p)$ has two branches, by considering the cut complex plane $\mathbb{C} \setminus \mathcal{N}$, it can be made analytic in this region. This implies (1). We now compute $S^\pm[\mathcal{F}](x)$:

$$S^\pm[\mathcal{F}](x) = \int_0^{e^{\pm i\theta}\infty} e^{-px} \mathcal{B}[\mathcal{F}_f](p) dp = C_M + \frac{3\pi c}{M^2 b} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) \int_0^{e^{\pm i\theta}\infty} \frac{e^{-px}}{\left(\frac{\ell^2\pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} dp \tag{2.10}$$

where the integral $\int_0^{e^{+i\theta}\infty}$ agrees with $\int_0^{+i\infty}$ if $\Im(x) > 0$ and the integral $\int_0^{e^{-i\theta}\infty}$ agrees with $\int_0^{-i\infty}$ if $\Im(x) < 0$. To prove the first part of (2), we start with the following formula, valid for

$\Re(x) > 0$:

$$\int_{\gamma} \frac{e^{-px}}{(1-p)^{\frac{5}{2}}} dp = -\frac{4}{3} + \frac{8}{3} \sqrt{\pi} \mathcal{E}(\sqrt{x}) \quad (2.11)$$

where $\gamma = [0, e^{i\theta}\infty) \cup [0, e^{-i\theta}\infty)$ and

$$\mathcal{E}(y) := \frac{2y^3 e^{-y^2}}{\sqrt{\pi}} \int_0^y e^{t^2} dt - \frac{y^2}{\sqrt{\pi}}.$$

From (1.3) and (2.10), we have

$$\begin{aligned} S_f^{\text{med}}(x) &= C_M + \frac{1}{2} \frac{3\pi c}{M^2 b} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) \int_{\gamma} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} dp \\ &= C_M + b \left(\frac{M}{\pi}\right)^3 \frac{3\pi c}{2M^2 b} \sum_{\ell=1}^{\infty} \frac{\tilde{f}(\ell)}{\ell} \int_{\gamma} \frac{e^{-\frac{b\ell^2 \pi^2 x}{M^2}}}{(1-p)^{5/2}} dp \end{aligned} \quad (2.12)$$

where we have made the change of variable $p \rightarrow \frac{b\ell^2 \pi^2}{M^2}$. Thus, (2.11) and (2.12) imply

$$\begin{aligned} S_f^{\text{med}}(x) &= C_M + \frac{3Mc}{2\pi^2} \sum_{\ell=1}^{\infty} \frac{\tilde{f}(\ell)}{\ell} \left\{ -\frac{4}{3} + \frac{8}{3} \sqrt{\pi} \mathcal{E}\left(\frac{\ell\pi}{M} \sqrt{bx}\right) \right\} \\ &= \frac{4Mc}{\pi^{3/2}} \sum_{\ell=1}^{\infty} \frac{\tilde{f}(\ell)}{\ell} \mathcal{E}\left(\frac{\ell\pi}{M} \sqrt{bx}\right) \end{aligned}$$

where one can check that

$$C_M = \frac{2Mc}{\pi^2} \sum_{\ell=1}^{\infty} \frac{\tilde{f}(\ell)}{\ell^2}. \quad (2.13)$$

This implies that $S_f^{\text{med}}(x)$ is analytic on $\Re(x) > 0$.

To prove the existence of the radial limit and part (3), we begin with rewriting the integral on the right-hand side of (2.10) as an integral of the partial theta function $\theta_{0,4M^2,f}^{(0)}(x)$. This requires some attention due to the branch cut singularity of the resulting square root. Integration by parts yields

$$\begin{aligned} \int_0^{e^{\pm i\theta}\infty} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} dp &= \left[\frac{2b}{3} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{3/2}} \right]_0^{e^{\pm i\theta}\infty} + \frac{2bx}{3} \int_0^{e^{\pm i\theta}\infty} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{3/2}} dp \\ &= \frac{2b}{3} \left[-\frac{M^3}{\ell^3 \pi^3} + x \int_0^{xe^{\pm i\theta}\infty} \frac{e^{-\frac{\ell^2 \pi^2 b}{M^2} p}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{\ell^2 \pi^2}{M^2 x} p\right)^{3/2}} d\left(\frac{\ell^2 \pi^2 b}{M^2 x} p\right) \right] \\ &= -\frac{2bM^3}{3\ell^3 \pi^3} + \frac{2b^2 M}{3\ell\pi} \int_0^{xe^{\pm i\theta}\infty} \frac{e^{-\frac{\ell^2 \pi^2 b}{M^2} p}}{\left(1 - \frac{p}{x}\right)^{3/2}} dp. \end{aligned} \quad (2.14)$$

The change of coordinate $x \rightarrow -\frac{1}{2\pi i\alpha}$ where $0 \neq \alpha \in \mathbb{C}$ to (2.14) yields

$$\begin{aligned} -\frac{2bM^3}{3\ell^3\pi^3} + \frac{2b^2M}{3\ell\pi} \int_0^{xe^{\pm i\theta}\infty} \frac{e^{-\frac{\ell^2\pi^2 b}{M^2}p}}{\left(1 - \frac{p}{x}\right)^{3/2}} dp &= -\frac{2bM^3}{3\ell^3\pi^3} + \frac{2b^2M}{3\ell\pi} \int_0^{-\frac{1}{2\pi i\alpha}e^{\pm i\theta}\infty} \frac{e^{2\pi i\frac{\ell^2 2\pi i b}{4M^2}p}}{(1 + 2\pi i p\alpha)^{3/2}} dp \\ &= -\frac{2bM^3}{3\ell^3\pi^3} + \frac{b^2M}{3i\ell\pi^2} \int_0^{-\frac{1}{\alpha}e^{\pm i\theta}\infty} \frac{e^{2\pi i\frac{\ell^2 b}{4M^2}p}}{(1 + p\alpha)^{3/2}} dp. \end{aligned} \quad (2.15)$$

Now $(1 + p\alpha)^{3/2}$ has a branch cut at $p = -1/\alpha$, hence we distinguish between $\Re(\alpha) > 0$ and $\Re(\alpha) < 0$ (see Figure 3). If $\Re(\alpha) > 0$, then $(1 + p\alpha)^{3/2} = \alpha^{3/2}(1/\alpha + p)^{3/2}$, and the integral at angle $-i\theta$ will not change sign as the path is not crossing the branch cut, i.e., $\int_0^{-\frac{1}{\alpha}e^{-i\theta}\infty} = \int_0^{+i\infty}$. If $\Re(\alpha) < 0$, then $(1 + p\alpha)^{3/2} = \alpha^{3/2}(1/\alpha + p)^{3/2}$, but the integral at angle $+i\theta$ will change sign as the path is crossing the branch cut, i.e., $\int_0^{-\frac{1}{\alpha}e^{+i\theta}\infty} = -\int_0^{+i\infty}$.




FIGURE 3. Branch cut of $(1 + p\alpha)^{3/2}$. On the left for $\Re(\alpha) < 0$ and on the right for $\Re(\alpha) > 0$.

By (2.10), (2.13) and (2.15), we have for $\alpha \in \mathbb{Q}_{>0}$

$$\begin{aligned} \int_0^{e^{-i\theta}\infty} e^{\frac{p}{2\pi i\alpha}} \mathcal{B}[\mathcal{F}_f](p) dp &= -i \frac{cb}{M\pi} \sum_{\ell=1}^{\infty} \tilde{f}(\ell) \int_0^{-\frac{1}{\alpha}e^{-i\theta}\infty} \frac{e^{2\pi i\frac{\ell^2 b}{4M^2}p}}{(1 + p\alpha)^{3/2}} dp \\ &= -i \frac{cb}{M\pi\alpha^{3/2}} \sum_{\ell=1}^{\infty} \tilde{f}(\ell) \int_0^{+i\infty} \frac{e^{2\pi i\frac{\ell^2 b}{4M^2}p}}{(1/\alpha + p)^{3/2}} dp \\ &= -i \frac{cb}{M\pi\alpha^{3/2}} \int_0^{+i\infty} \frac{\theta^{(0)}_{0,4M^2,\tilde{f}}(bp)}{(1/\alpha + p)^{3/2}} dp \end{aligned}$$

while for $\alpha \in \mathbb{Q}_{<0}$, we obtain

$$\int_0^{e^{+i\theta}\infty} e^{\frac{p}{2\pi i\alpha}} \mathcal{B}[\mathcal{F}_f](p) dp = -i \frac{cb}{M\pi} \sum_{\ell=1}^{\infty} \tilde{f}(\ell) \int_0^{-\frac{1}{\alpha}e^{+i\theta}\infty} \frac{e^{2\pi i\frac{\ell^2 b}{4M^2}p}}{(1 + p\alpha)^{3/2}} dp$$

$$\begin{aligned}
&= i \frac{cb}{M\pi\alpha^{3/2}} \sum_{\ell=1}^{\infty} \tilde{f}(\ell) \int_0^{+i\infty} \frac{e^{2\pi i \frac{\ell^2 b}{4M^2} p}}{(1/\alpha + p)^{3/2}} dp \\
&= i \frac{cb}{M\pi\alpha^{3/2}} \int_0^{+i\infty} \frac{\theta_{0,4M^2, \tilde{f}}^{(0)}(bp)}{(1/\alpha + p)^{3/2}} dp.
\end{aligned}$$

We now compute the discontinuity $\text{disc}^0(x)$ across the positive real axis:


$$\begin{aligned}
\text{disc}^0(x) &:= S^+[\mathcal{F}](x) - S^-[\mathcal{F}](x) \\
&= \frac{3\pi c}{M^2 b} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) \left[\int_0^{e^{+i\theta}\infty} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} dp - \int_0^{e^{-i\theta}\infty} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} dp \right] \\
&= \frac{3\pi c}{M^2 b} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) \int_{\mathcal{C}} \frac{e^{-px}}{\left(\frac{\ell^2 \pi^2}{M^2} - \frac{p}{b}\right)^{5/2}} dp \\
&= \frac{3\pi c b^{3/2}}{M^2} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) e^{-\frac{\ell^2 \pi^2 b}{M^2} x} \int_{\mathcal{C}_\ell} \frac{e^{-px}}{(-p)^{5/2}} dp \\
&= \frac{3\pi c b^{3/2}}{M^2} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) e^{-\frac{\ell^2 \pi^2 b}{M^2} x} \int_{\mathcal{C}_\ell} \frac{e^{-px}}{(-xp)^{5/2}} d(xp) x^{3/2} \\
&= \frac{2\pi i}{\Gamma\left(\frac{5}{2}\right)} x^{3/2} \frac{3\pi c b^{3/2}}{M^2} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) e^{-\frac{\ell^2 \pi^2 b}{M^2} x} \\
&= 2i (2b\pi x)^{3/2} \frac{\sqrt{2}c}{M^2} \sum_{\ell=1}^{\infty} \ell \tilde{f}(\ell) e^{-\frac{\ell^2 \pi^2 b}{M^2} x} \\
&= 2i (2b\pi x)^{3/2} \frac{\sqrt{2}c}{M^2} \theta_{0,4M^2, \tilde{f}}^{(1)}(2\pi i x).
\end{aligned} \tag{2.16}$$

Here, $\mathcal{C}_\ell = \mathcal{C} - \frac{\ell^2 \pi^2 b}{M^2}$ where \mathcal{C} is the Hankel contour enclosing the branch point singularity arising from the second line in (2.16), as in Figure 4. The radial limit $x \rightarrow -\frac{1}{2\pi i \alpha}$ of the partial theta series appearing in the last line of (2.16) exists by [2, Proposition 3.1] and so

$$\text{disc}^0\left(-\frac{1}{2\pi i \alpha}\right) = 2i \left(\frac{ib}{\alpha}\right)^{3/2} \frac{\sqrt{2}c}{M^2} \theta_{0,4M^2, \tilde{f}}^{(1)}\left(-\frac{b}{\alpha}\right).$$

In addition, the radial limit $x \rightarrow -\frac{1}{2\pi i \alpha}$ also exists for $S^\pm[\mathcal{F}](x)$ since they are analytic functions on the half-planes $\Re(e^{\pm i\theta} x) > 0$, respectively (see the change of coordinates after (2.14)). Thus, the second part of (2) follows. Summarizing, we find that

$$S_f^{\text{med}}\left(-\frac{1}{2\pi i \alpha}\right) = \begin{cases} S^+\left(-\frac{1}{2\pi i \alpha}\right) + \left(\frac{b}{\alpha}\right)^{3/2} e^{\pi i/4} \frac{\sqrt{2}c}{M^2} \theta_{0,4M^2, \tilde{f}}^{(1)}\left(-\frac{b}{\alpha}\right) & \text{if } \alpha \in \mathbb{Q}_{>0}, \\ S^-\left(-\frac{1}{2\pi i \alpha}\right) - \left(\frac{b}{\alpha}\right)^{3/2} e^{\pi i/4} \frac{\sqrt{2}c}{M^2} \theta_{0,4M^2, \tilde{f}}^{(1)}\left(-\frac{b}{\alpha}\right) & \text{if } \alpha \in \mathbb{Q}_{<0} \end{cases}$$

FIGURE 4. The Hankel contour \mathcal{C} for the discontinuity $\text{disc}^0(x)$.

with

$$S^\pm \left(-\frac{1}{2\pi i \alpha} \right) = \mp i \frac{cb}{M\pi\alpha^{3/2}} \int_0^{+i\infty} \frac{\theta_{0,4M^2, \tilde{f}}^{(0)}(bp)}{(1/\alpha + p)^{3/2}} dp.$$

Finally, noticing that $(i\alpha)^{-3/2} = \mp e^{\pi i/4} \alpha^{-3/2}$ depending on the sign of α and on our choice of the branch of the square root, part (3) follows. \square

3. PROOF OF THEOREM 1.4 AND COROLLARY 1.5

In order to prove Theorem 1.4 and Corollary 1.5, we require some general preliminaries. Recall the periodic function $\chi_{2st}^{(n,m)}(k)$ given by (1.18).

Lemma 3.1. *For all $k \in \mathbb{Z}$, $1 \leq m \leq t-1$ and $1 \leq n \leq s-1$, we have*

$$\chi_{2st}^{(n,m)}(k) = \chi_{2st}^{(s-n, t-m)}(k).$$

Proof. By replacing $n \rightarrow s-n$ and $m \rightarrow t-m$, we see that

$$\begin{aligned} \pm((s-n)t - (t-m)s) &= \pm(ms - nt) = \mp(nt - ms), \\ \pm((s-n)t + (t-m)s) &= \pm(2st - ms - nt) \equiv \mp(nt + ms) \pmod{2st}, \end{aligned} \quad (3.1)$$

and thus the result follows from (3.1). \square

In view of Lemma 3.1, the map

$$(n, m) \rightarrow \begin{cases} (n, m) & \text{if } 1 \leq n \leq \frac{s-1}{2}, \quad 1 \leq m \leq \frac{t-1}{2}, \\ (s-n, t-m) & \text{otherwise.} \end{cases} \quad (3.2)$$

is a bijection between the pairs defined by $\mathcal{D}_1(s, t)$ and $\mathcal{D}_2(s, t)$ in (1.19) and (1.20), respectively.

Lemma 3.2. *Let s and t be coprime positive integers and $\mathcal{D}(s, t)$ be the set given by (1.21). Then the integers in the set*

$$\mathcal{S} = \{\pm(nt \pm ms) : (n, m) \in \mathcal{D}(s, t)\}$$

are distinct and thus $|\mathcal{S}| = 2(s-1)(t-1)$.

Proof. Without loss of generality, we can assume that s and t are coprime odd integers. Let us consider two integers of the forms $n_1t + m_1s$ and $n_2t + m_2s$ where $-\frac{s-1}{2} \leq n_1, n_2 \leq \frac{s-1}{2}$ and $-(t-1) \leq m_1, m_2 \leq t-1$. Then

$$n_1t + m_1s = n_2t + m_2s \iff (n_1 - n_2)t + (m_1 - m_2)s = 0. \quad (3.3)$$

Since s and t are coprime, (3.3) implies that $s \mid (n_1 - n_2)$, which yields $|n_1 - n_2| \geq s$. On the other hand, since $-\frac{s-1}{2} \leq n_1, n_2 \leq \frac{s-1}{2}$, we obtain $-(s-1) \leq n_1 - n_2 \leq s-1$, or equivalently, $|n_1 - n_2| \leq s-1$, a contradiction. This shows that $n_1 = n_2$ and thus (3.3) implies that $m_1 = m_2$. \square

Define

$$S_{n,m}^{n',m'} = \sqrt{\frac{8}{st}} (-1)^{nm'+mn'+1} \sin\left(\frac{nn't}{s}\pi\right) \sin\left(\frac{mm's}{t}\pi\right).$$

Proposition 3.3. *For all $k \in \mathbb{Z}$, and $(n, m) \in \mathcal{D}(s, t)$, we have*

$$\tilde{\chi}_{2st}^{(n,m)}(k) = -\sqrt{\frac{st}{8}} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n,m}^{n',m'} \chi_{2st}^{(n',m')}(k). \quad (3.4)$$

Proof. By (1.12), we have

$$\begin{aligned} \tilde{\chi}_{2st}^{(n,m)}(k) &= (-1)^k \sin\left(\frac{(nt + ms - (nt - ms))k\pi}{2st}\right) \sin\left(\frac{(2st - (nt + ms) - (nt - ms))k\pi}{2st}\right) \\ &= (-1)^k \sin\left(\frac{mk\pi}{t}\right) \sin\left(\frac{(s-n)k\pi}{s}\right) \\ &= -\sin\left(\frac{mk\pi}{t}\right) \sin\left(\frac{nk\pi}{s}\right). \end{aligned} \quad (3.5)$$

Clearly, the support of $\tilde{\chi}_{2st}^{(n,m)}$ is contained within the set of integers k such that $s \nmid k$ and $t \nmid k$. By inclusion-exclusion, the total number of integers between 0 and $2st - 1$ which are either multiples of s or t is $2s + 2t - 2$. By Lemma 3.2, the number of distinct integers $k = \pm(m's \pm n't)$ between 0 and $2st - 1$, where $(n', m') \in \mathcal{D}(s, t)$ is $2(s-1)(t-1)$. Since $\tilde{\chi}_{2st}^{(n,m)}$ has period $2st$, this implies that the support of $\tilde{\chi}_{2st}^{(n,m)}$ is contained in the following set of distinct integers

$$\mathcal{T} = \{\pm(m's \pm n't) \pmod{2st} : (n', m') \in \mathcal{D}(s, t)\}. \quad (3.6)$$

Thus it suffices to prove (3.4) for integers in \mathcal{T} . From (3.5), we have

$$\begin{aligned} \tilde{\chi}_{2st}^{(n,m)}(\pm(m's \pm n't)) &= -\sin\left(\frac{\pm m(m's \pm n't)\pi}{t}\right) \sin\left(\frac{\pm n(m's \pm n't)\pi}{s}\right) \\ &= -\sin\left(\pm mn'\pi + \frac{mm's}{t}\pi\right) \sin\left(nm'\pi \pm \frac{nn't}{s}\pi\right) \\ &= \pm(-1)^{mn'+nm'+1} \sin\left(\frac{mm's}{t}\pi\right) \sin\left(\frac{nn't}{s}\pi\right) \\ &= (-1)^{mn'+nm'} \sin\left(\frac{mm's}{t}\pi\right) \sin\left(\frac{nn't}{s}\pi\right) \chi_{2st}^{(n',m')}(\pm(m's \pm n't)) \end{aligned} \quad (3.7)$$

where in the last step we use that the sign \pm depends on the sign of $(m's \pm n't)$, and it is $-\chi_{2st}^{(n',m')}(\pm(m's \pm n't))$. Since the integers in \mathcal{T} are distinct, then from (3.7) we find:

$$\tilde{\chi}_{2st}^{(n,m)}(\pm(m's \pm n't)) = -\sqrt{\frac{st}{8}} \sum_{(n'',m'') \in \mathcal{D}(s,t)} S_{n'',m''}^{n',m'} \chi_{2st}^{(n'',m'')}(\pm(m's \pm n't)). \quad (3.8)$$

It follows from (3.8) that (3.4) is true for integers k given by (3.6), which are non-multiples of s or t and between 0 and $2st - 1$. It remains to show that (3.4) remains true when k is a multiple of s or t . Without loss of generality, we assume that k is a multiple of s . Then (3.5) implies that $\tilde{\chi}_{2st}^{(n,m)}(k) = 0$. Thus, it suffices to show that $\chi_{2st}^{(n',m')}(k) = 0$ for all $(n',m') \in \mathcal{D}(s,t)$. To this end, we note that since s and t are coprime, $s \mid \pm(n't \pm m's)$ if and only if $s \mid n'$, which implies that $n' \geq s$, a contradiction. Thus, for $(n',m') \in \mathcal{D}(s,t)$, it follows $\pm(n't \pm m's) \not\equiv 0 \pmod{s}$. Hence $k \neq \pm(n't \pm m's)$, and $\chi_{2st}^{(n',m')}(k) = 0$. \square

Finally, we recall two key properties for the non-holomorphic Eichler integral

$$\widehat{\Phi}^{(n,m)}(z) := \sqrt{\frac{sti}{8\pi^2}} \int_{\bar{z}}^{+i\infty} \frac{\theta_{0,4st,\chi_{2st}}^{(n,m)}(\tau)}{(\tau - z)^{3/2}} d\tau$$

where $z \in \mathbb{H}^-$ [37, Eqn. (16)]. For $z \in \mathbb{H}^-$ and $\alpha \in \mathbb{Q}$, consider the period function

$$r^{(n,m)}(z; \alpha) := \sqrt{\frac{sti}{8\pi^2}} \int_{\alpha}^{+i\infty} \frac{\theta_{0,4st,\chi_{2st}}^{(n,m)}(\tau)}{(\tau - z)^{3/2}} d\tau,$$

as defined in [37, Eqn. (18)]. Then, we have

$$\widehat{\Phi}^{(n,m)}(z) + \left(\frac{1}{iz}\right)^{3/2} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n',m'}^{n',m'} \widehat{\Phi}^{(n',m')} \left(-\frac{1}{z}\right) = r^{(n,m)}(z; 0) \quad (3.9)$$

and

$$\widehat{\Phi}^{(n,m)}(\alpha) = -\frac{1}{2} \theta_{0,4st,\chi_{2st}}^{(1)}(\alpha) \quad (3.10)$$

where (3.9) is [37, Eqn. (17)] and (3.10) is [37, below Eqn. (19)].

Proof of Theorem 1.4. By Proposition 3.3, we have

$$\theta_{0,4(2st)^2,\tilde{\chi}_{2st}^{(n,m)}}^{(\nu)}(4stz) = -\sqrt{\frac{st}{8}} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n',m'}^{n',m'} \theta_{0,4st,\chi_{2st}}^{(\nu)}(z). \quad (3.11)$$

Also, [37, Eqn. (10)] states

$$\theta_{0,4st,\chi_{2st}}^{(0)}(z) = \sqrt{\frac{i}{z}} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n',m'}^{n',m'} \theta_{0,4st,\chi_{2st}}^{(0)} \left(-\frac{1}{z}\right). \quad (3.12)$$

Applying Theorem 1.2 part (3) and (3.11) gives

$$S_{\chi_{2st}^{(n,m)}}^{\text{med}} \left(-\frac{1}{2\pi i\alpha}\right) = -\frac{2e^{\pi i/4}}{\pi(i\alpha)^{3/2}} \sqrt{\frac{st}{8}} \int_0^{+i\infty} \frac{\sum_{(n',m') \in \mathcal{D}(s,t)} S_{n',m'}^{n',m'} \theta_{0,4st,\chi_{2st}}^{(0)}(p)}{\left(\frac{1}{\alpha} + p\right)^{3/2}} dp$$

$$- \frac{1}{(i\alpha)^{3/2}} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n,m}^{n',m'} \theta_{0,4st,\chi_{2st}^{(n',m')}}^{(1)} \left(-\frac{1}{\alpha} \right). \quad (3.13)$$

Then, by (3.12), we rewrite (3.13) as follows:

$$\begin{aligned} S_{\chi_{2st}^{(n,m)}}^{\text{med}} \left(-\frac{1}{2\pi i\alpha} \right) &= -\frac{2e^{\pi i/4}}{\pi(i\alpha)^{3/2}} \sqrt{\frac{sti}{8}} \int_0^{+i\infty} \frac{\theta_{0,4st,\chi_{2st}^{(n,m)}}^{(0)}(-1/p)}{\sqrt{p} \left(\frac{1}{\alpha} + p \right)^{3/2}} dp \\ &\quad - (i\alpha)^{-3/2} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n,m}^{n',m'} \theta_{0,4st,\chi_{2st}^{(n',m')}}^{(1)} \left(-\frac{1}{\alpha} \right). \end{aligned} \quad (3.14)$$

We now perform the change of coordinates $p \rightarrow -1/p$ to the integral in (3.14) and simplify to obtain

$$\frac{2e^{\pi i/4}}{\pi(i\alpha)^{3/2}} \sqrt{\frac{sti}{8}} \int_0^{+i\infty} \frac{\theta_{0,4st,\chi_{2st}^{(n,m)}}^{(0)}(-1/p)}{\sqrt{p} \left(\frac{1}{\alpha} + p \right)^{3/2}} dp = 2\sqrt{\frac{sti}{8\pi^2}} \int_0^{+i\infty} \frac{\theta_{0,4st,\chi_{2st}^{(n,m)}}^{(0)}(p)}{(p-\alpha)^{3/2}} dp.$$

Then, going back to (3.14) we find

$$\begin{aligned} S_{\chi_{2st}^{(n,m)}}^{\text{med}} \left(-\frac{1}{2\pi i\alpha} \right) &= -2\sqrt{\frac{sti}{8\pi^2}} \int_0^{+i\infty} \frac{\theta_{0,4st,\chi_{2st}^{(n,m)}}^{(0)}(p)}{(p-\alpha)^{3/2}} dp \\ &\quad - (i\alpha)^{-3/2} \sum_{(n',m') \in \mathcal{D}(s,t)} S_{n,m}^{n',m'} \theta_{0,4st,\chi_{2st}^{(n',m')}}^{(1)} \left(-\frac{1}{\alpha} \right) \\ &= \theta_{0,4st,\chi_{2st}^{(n,m)}}^{(1)}(\alpha) \end{aligned} \quad (3.15)$$

where the last line of (3.15) follows from (3.9) and (3.10). This proves the result. \square

Proof of Corollary 1.5. Let $u \in \mathbb{Z}_{\geq 1}$. For $0 \leq \ell \leq u-1$, define

$$X_u^{(\ell)}(q) := \sum_{k_1, k_2, \dots, k_u=0}^{\infty} (q)_{k_u} q^{k_1^2 + \dots + k_{u-1}^2 + k_{\ell+1} + \dots + k_{u-1}} \prod_{i=1}^{u-1} \begin{bmatrix} k_{i+1} + \delta_{i,\ell} \\ k_i \end{bmatrix} \quad (3.16)$$

where $\delta_{i,\ell}$ is the characteristic function; see [36, Eqn. (11)]. The expression $X_u^{(\ell)}(q)$ matches the N th colored Jones polynomial for $T(2, 2u+1)$ when $\ell = 0$ and $q = \zeta_N$ and is an element of \mathcal{H} ; see [36, Proposition 16]. Take $s = 2$, $t = 2u+1$, $n = 1$ and $m = \ell+1$ in (1.18). Then Hikami's strange identity [36, Eqn. (15)]¹ reads

$$X_u^{(\ell)}(q) \text{ " = " } - \frac{1}{2} \theta_{(2u-2\ell-1)^2, 2(8u+4), \chi_{4(2u+1)}^{(1,\ell+1)}}^{(1)}(q). \quad (3.17)$$

Claims (1) and (2) of Conjecture 1.1 follow from Theorem 1.2 parts (1) and (2), namely the formal series $\mathcal{F}_{\chi_{4(2u+1)}^{(1,\ell+1)}}(x)$ has a resurgent Borel transform and its median resummation is an

¹Taking $u = 1$ and $\ell = 0$ in (3.16) and (3.17) yields (1.6).

analytic function for $\Re(x) > 0$ with radial limits at points $\frac{1}{2\pi i}\mathbb{Q}$. In particular, this is true for

$$F_{T(2,2u+1)}(x) = \mathcal{F}_{\chi_{4(2u+1)}^{(1,1)}}(x).$$

Then, part (3) of Conjecture 1.1 follows by Theorem 1.4 applying (3.17) for $\ell = 0$. In fact, we can prove a stronger result:

$$S_{\chi_{4(2u+1)}^{(1,\ell+1)}}^{\text{med}} \left(-\frac{1}{2\pi i\alpha} \right) \doteq \mathcal{F}_{\chi_{4(2u+1)}^{(1,\ell+1)}}(e^{2\pi i\alpha}) \quad (3.18)$$

for every $0 \neq \alpha \in \mathbb{Q}$ and $0 \leq \ell \leq u - 1$.

Let $k \in \mathbb{Z}_{\geq 1}$ and $\mathfrak{F}_k(q)$ denote the element in \mathcal{H} that matches the N th colored Jones polynomial for $T(3, 2^k)$ at a root of unity $q = e^{\frac{2\pi i}{N}}$ (see [6, Eqn. (1.8)] for an explicit q -hypergeometric expression). Choose $s = 3$, $t = 2^k$, $n = 2$ and $m = 1$ in (1.18). Then the strange identity

$$\mathfrak{F}_k(q) \text{ “} = \text{”} - \frac{1}{2} \theta_{(2^{k+1}-3)^2, 3 \cdot 2^{k+2}, \chi_{3 \cdot 2^{k+1}}^{(2,1)}}^{(1)}(q) \quad (3.19)$$

was proved in [6, Theorem 2.4]². Claims (1) and (2) of Conjecture 1.1 follow from Theorem 1.2 parts (1) and (2), namely the formal series

$$F_{T(3,2^k)}(x) = \mathcal{F}_{\chi_{3 \cdot 2^{k+1}}^{(2,1)}}(x)$$

has a resurgent Borel transform and its median resummation is an analytic function for $\Re(x) > 0$ with radial limits at points $\frac{1}{2\pi i}\mathbb{Q}$. Then, part (3) of Conjecture 1.1 follows by Theorem 1.4 applying (3.19). \square

4. CONCLUDING REMARKS

In order to apply Theorems 1.2 and 1.4 and thus verify Conjecture 1.1 for all torus knots $T(s, t)$, one needs to prove the relevant strange identity in (1.13). The first obstruction in this task is finding an explicit “non-cyclotomic” expansion for $J_N(T(s, t); q)$ from which an element in \mathcal{H} such as the one in (3.16) can be extracted. The Rosso-Jones formula for $J_N(T(s, t); q)$ does not appear to be sufficient [50, page 132]. Instead, one should consider the walks along braids method in [5, 7]. The second obstruction is in determining the underlying q -series identity which implies the strange identity. Thus, it would be of substantial interest to further develop the techniques in [45]. Finally, what can one say about Conjecture 1.1 for satellite or hyperbolic knots?

ACKNOWLEDGEMENTS

The first author thanks Dan Zhang, Nick Dorey and Phil Trinh for many interesting discussions. He was supported by EPSRC grant EP/V012479/1 as a postdoc at the University of Bath while part of this work was completed. The fourth author was partially supported by Enterprise Ireland CS20212030 and the Irish Research Council Advanced Laureate Award IRCLA/2023/1934. He would like to thank the Okinawa Institute of Science and Technology for their hospitality and support during his visit from June 9 to September 1, 2022 as part of their Theoretical Sciences Visiting Program. In particular, the first and fourth authors are very grateful to Professor Reiko Toriumi (OIST) for her firm instance for us to converse after the

²Taking $k = 1$ in (3.19) gives (1.6).

first author’s talk at OIST on August 23, 2022. The fourth author would also like to thank the Max-Planck-Institut für Mathematik for their hospitality and support during the completion of this paper. The fifth author has been supported by the Huawei Young Talents Program at IHÉS. Finally, the authors thank David Sauzin for his helpful comments on median resummation and the referees for their critiques which substantially improved the paper.

REFERENCES

- [1] S. Ahlgren, B. Kim, *Dissections of a “strange” function*, Int. J. Number Theory **11** (2015), no. 5, 1557–1562.
- [2] S. Ahlgren, B. Kim and J. Lovejoy, *Dissections of strange q -series*, Ann. Comb. **23** (2019), no. 3-4, 427–442.
- [3] J. Andersen, W. Mistegård, *Resurgence analysis of quantum invariants of Seifert fibered homology spheres*, J. Lond. Math. Soc. (2) **105** (2022), no. 2, 709–764.
- [4] G. E. Andrews, J. Sellers, *Congruences for the Fishburn numbers*, J. Number Theory **161** (2016), 298–310.
- [5] C. Armond, *Walks along braids and the colored Jones polynomial*, J. Knot Theory Ramifications **23** (2014), no. 2, 1450007, 15pp.
- [6] C. Bijaoui, H.U. Boden, B. Myers, R. Osburn, W. Rushworth, A. Tronsgard and S. Zhou, *Generalized Fishburn numbers and torus knots*, J. Combin. Theory Ser. A **178** (2021), 105355.
- [7] H.U. Boden, M. Shimoda, *Braid representatives minimizing the number of simple walks*, Ars Math. Contemp. **23** (2023), no. 1, Paper No. 10, 27 pp.
- [8] M. Borinsky, G. V. Dunne, *Non-perturbative completion of Hopf-algebraic Dyson-Schwinger equations*, Nuclear Phys. B **957** (2020), 115096, 17 pp.
- [9] F. Chapoton, F. Fauvet, C. Malvenuto and J-Y, Thibon, *Algebraic combinatorics, resurgence, moulds and applications (CARMA). Volume 1*, IRMA Lectures in Mathematics and Theoretical Physics, **31**. EMS Publishing House, Berlin 2020.
- [10] F. Chapoton, F. Fauvet, C. Malvenuto and J-Y, Thibon, *Algebraic combinatorics, resurgence, moulds and applications (CARMA). Volume 2*, IRMA Lectures in Mathematics and Theoretical Physics, **32**. EMS Publishing House, Berlin 2020.
- [11] O. Costin, *On Borel summation and Stokes phenomena for rank-1 nonlinear systems of ordinary differential equations*, Duke Math. J. **93** (1998), no. 2, 289–344.
- [12] O. Costin, *Asymptotics and Borel summability*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, **141** CRC Press, Boca Raton, FL, 2009.
- [13] O. Costin, S. Garoufalidis, *Resurgence of the Kontsevich-Zagier series*, Ann. Inst. Fourier (Grenoble) **61** (2011), no. 3, 1225–1258.
- [14] E. Delabaere, F. Pham, *Resurgent methods in semi-classical asymptotics*, Ann. Inst. H. Poincaré Phys. Théor. **71** (1999), no. 1, 1–94.
- [15] D. Dorigoni, A. Kleinschmidt, *Resurgent expansion of Lambert series and iterated Eisenstein integrals*, Commun. Number Theory Phys. **15** (2021), no. 1, 1–57.
- [16] J. Écalle, *Les fonctions résurgentes. Tome I. Les algèbres de fonctions résurgentes*, Publications Mathématiques d’Orsay, vol. 81, p. 5, Université de Paris-Sud, Département de Mathématique, Orsay, 1981, 247 pp.
- [17] J. Écalle, *Les fonctions résurgentes. Tome II. Les fonctions résurgentes appliquées à l’itération*, Publications Mathématiques d’Orsay, vol. 81, p. 6, Université de Paris-Sud, Département de Mathématique, Orsay, 1981, pp. 248–531.
- [18] J. Écalle, *Les fonctions résurgentes. Tome III. L’équation du pont et la classification analytique des objets locaux*, Publications Mathématiques d’Orsay, vol. 85, p. 5, Université de Paris-Sud, Département de Mathématique, Orsay, 1985, 587 pp.
- [19] V. Fantini, C. Wheeler, *Summability for state integrals of hyperbolic knots*, preprint available at [arXiv:2410.20973](https://arxiv.org/abs/2410.20973).
- [20] S. Garoufalidis, *On the characteristic and deformation varieties of a knot*, Proceedings of the Casson Fest, 291–309, Geom. Topol. Monogr., **7**, Geom. Topol. Publ., Coventry, 2004.
- [21] S. Garoufalidis, *Chern-Simons theory, analytic continuation and arithmetic*, Acta Math. Vietnam. **33** (2008), no. 3, 335–362.

- [22] S. Garoufalidis, *The Jones slopes of a knot*, Quantum Topol. **2** (2011), no. 1, 43–69.
- [23] S. Garoufalidis, T. T. Q. Lê, *Gevrey series in quantum topology*, J. Reine Angew. Math. **618** (2008), 169–195.
- [24] S. Garoufalidis, J. Gu and M. Mariño, *The resurgent structure of quantum knot invariants*, Comm. Math. Phys. **386** (2021), no. 1, 469–493.
- [25] S. Garoufalidis, J. Gu and M. Mariño, *Peacock patterns and resurgence in complex Chern-Simons theory*, Res. Math. Sci. **10** (2023), no. 3, Paper No. 29, 67 pp.
- [26] S. Garoufalidis, J. Gu, M. Mariño and C. Wheeler, *Resurgence of Chern-Simons theory at the trivial flat connection*, preprint available at [arXiv:2111.04763](https://arxiv.org/abs/2111.04763).
- [27] A. Goswami, *Congruences for generalized Fishburn numbers at roots of unity*, Acta Arith. **199** (2021), no. 1, 77–102.
- [28] A. Goswami, A. Jha, B. Kim and R. Osburn, *Asymptotics and sign patterns for coefficients in expansions of Habiro elements*, Math. Z., **304** (2023), no. 4, Paper No. 57, pp. 17.
- [29] A. Goswami, R. Osburn, *Quantum modularity of partial theta series with periodic coefficients*, Forum Math. **33** (2021), no. 2, 451–463.
- [30] K. Habiro, *On the colored Jones polynomial of some simple links*, in: Recent progress toward the volume conjecture (Kyoto, 2000), Sūrikaiseikikenkyūsho Kōkyūroku **1172** (2000), 34–43.
- [31] K. Habiro, *Cyclotomic completions of polynomials rings*, Publ. Res. Inst. Math. Sci. **40** (2004), no. 4, 1127–1146.
- [32] L. Han, Y. Li, D. Sauzin and S. Sun, *Resurgence and partial theta series*, Funct. Anal. Appl. **57** (2023), no. 3, 248–265.
- [33] K. Hikami, *On the quantum invariant for the Brieskorn homology spheres*, Internat. J. Math. **16** (2005), no. 6, 661–685.
- [34] K. Hikami, *Quantum invariant, modular form, and lattice points*, Int. Math. Res. Not. (2005), no. 3, 121–154.
- [35] K. Hikami, *Quantum invariants, modular forms, and lattice points. II*, J. Math. Phys. **47** (2006), no. 10, 102301, 32 pp.
- [36] K. Hikami, *q-series and L-functions related to half-derivates of the Andrews-Gordon identity*, Ramanujan J. **11** (2006), no. 2, 175–197.
- [37] K. Hikami, A.N. Kirillov, *Torus knot and minimal model*, Phys. Lett. B **575** (2003), no. 3-4, 343–348.
- [38] V. Huynh, T. T. Q. Lê, *On the colored Jones polynomial and the Kashaev invariant*, J. Math. Sci. (N.Y.) **146** (2007), no. 1, 5490–5504.
- [39] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc. (N.S.) **12** (1985), no. 1, 103–111.
- [40] E. Kalfagianni, A. T. Tran, *Knot cabling and the degree of the colored Jones polynomial*, New York J. Math. **21** (2015), 905–941.
- [41] R. Kashaev, *Quantum dilogarithm as a 6j-symbol*, Modern Phys. Lett. A **9** (1994), no. 40, 3757–3768.
- [42] S. Kamimoto, *Resurgent functions and nonlinear systems of differential and difference equations*, Adv. Math. **406** (2022), Paper No. 108533, 28pp.
- [43] M. Kontsevich, Y. Soibelman, *Analyticity and resurgence in wall-crossing formulas*, Lett. Math. Phys. **112** (2022), no. 2, Paper No. 32, 56 pp.
- [44] T. T. Q. Lê, *Quantum invariants of 3-manifolds: Integrality, splitting, and perturbative expansion*, Topology Appl. **127** (2003), no. 1-2, 125–152.
- [45] J. Lovejoy, *Bailey pairs and strange identities*, J. Korean Math. Soc. **59** (2022), no. 5, 1015–1045.
- [46] A. A. Mahmoud, K. Yeats, *Connected chord diagrams and the combinatorics of asymptotic expansions*, J. Integer Seq. **22** (2022), no. 7, Art. 22.7.5, 22 pp.
- [47] B. Malgrange, *Introduction aux travaux de J. Écalle*, Enseign. Math. (2) **31** (1985), no. 3-4, 261–282.
- [48] M. Mariño, *Lectures on non-perturbative effects in large N gauge theories, matrix models and strings*, Fortschr. Phys. **62** (2014), no. 5-6, 455–540.
- [49] C. Mitschi, D. Sauzin, *Divergent series, summability and resurgence. I. Monodromy and resurgence*, Lecture Notes in Mathematics, **2153** Springer, 2016.
- [50] H.R. Morton, *The coloured Jones function and Alexander polynomial for torus knots*, Math. Proc. Cambridge Philos. Soc. **117** (1995), no. 1, 129–135.
- [51] H. Murakami, Y. Yokota, *Volume conjecture for knots*, SpringerBriefs in Mathematical Physics, 30. Springer, Singapore, 2018.

- [52] D. Sauzin, *Variations on the resurgence of the Gamma function*, preprint available at [arXiv:2112.15226](https://arxiv.org/abs/2112.15226).
- [53] A. Straub, *Congruences for Fishburn numbers modulo prime powers*, *Int. J. Number Theory* **11** (2015), no. 5, 1679–1690.
- [54] D. Zagier, *Vassiliev invariants and a strange identity related to the Dedekind eta-function*, *Topology* **40** (2001), no. 5, 945–960.
- [55] D. Zagier, *Quantum modular forms*, *Quanta of maths*, 659–675, *Clay Math. Proc.*, **11**, Amer. Math. Soc., Providence, RI, 2010.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, SOUTH KENSINGTON CAMPUS, LONDON SW7 2AZ, UK

Email address: `samuel.crew24@imperial.ac.uk`

LABORATOIRE MATHÉMATIQUE ORSAY, BUILDING 307, RUE MICHEL MAGAT, FACULTÉ DES SCIENCES D’ORSAY, UNIVERSITÉ PARIS-SACLAY, F-91405 ORSAY CEDEX, FRANCE

Email address: `veronica.fantini@universite-paris-saclay.fr`

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, THE UNIVERSITY OF TEXAS RIO GRANDE VALLEY, 1201 W. UNIVERSITY DR., EDINBURG, TX, 78539

Email address: `ankush.goswami@utrgv.edu`

Email address: `ankushgoswami3@gmail.com`

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY COLLEGE DUBLIN, BELFIELD, DUBLIN 4, IRELAND

Email address: `robert.osburn@ucd.ie`

THEORETICAL SCIENCES VISITING PROGRAM, OKINAWA INSTITUTE OF SCIENCE AND TECHNOLOGY GRADUATE UNIVERSITY, ONNA, 904-0495, JAPAN

IHÉS, LE BOIS-MARIE, 35, ROUTE DE CHARTRES, 91440 BURES-SUR-YVETTE, FRANCE

Email address: `wheeler@ihes.fr`