

# Gravitational Modelling Summer School, Dublin 2018: Astrophysics and Data Analysis Workshop

Jonathan R. Gair

*School of Mathematics, University of Edinburgh, James Clerk Maxwell Building,  
King's Buildings, Peter Guthrie Tait Road, Edinburgh, EH9 3FD\**

## I. INTRODUCTION

This workshop has two parts. The first part is a set of calculations based on the material in lecture 1 of this course. For two different classes of source you will estimate properties such as the characteristic strains, signal-to-noise ratios and background spectrum generated by a population of such sources. There are two sub-parts to this question. At the end of each there is a starred part which is more open ended and involves some numerical investigation. I suggest you leave that to the end of or after the workshop.

The second part of the lecture is a computing practical which will show you how to use freely available python packages to carry out some of the parameter estimation and hierarchical modelling techniques described in the second half of the third lecture. There are no questions as such — all the necessary code is included in the python notebook. However, it is worth working through that material carefully to check it all makes sense and do try playing around with the code to modify some of the properties of the model.

## II. PART I: COMPUTING GW BACKGROUND SPECTRA

### A. Massive black hole binary background

This question is concerned with properties of a background generated by a population of massive black holes. For the first few parts of this question you can assume that these binaries are identical, i.e., they all have the same values for the two masses,  $m_1$  and  $m_2$ , and hence the derived quantities of total mass,  $M = m_1 + m_2$ , reduced mass,  $\mu = m_1 m_2 / M$ , and chirp mass,

$$\mathcal{M}_c = \frac{m_1^{\frac{3}{5}} m_2^{\frac{3}{5}}}{M^{\frac{1}{5}}}.$$

In the final part of this question you'll be asked to think about and play around with different assumptions about the mass distribution in the binaries.

We will use geometric units throughout, i.e., we set  $c = G = 1$  so we don't need to worry about keeping track of these factors.

(1) Assuming that the binary is Newtonian and circular, reproduce the various scaling relations derived in lecture 1, in particular show that

(i) The GW amplitude scales like

$$h \sim \frac{1}{D} \mathcal{M}_c^{\frac{5}{3}} f^{\frac{2}{3}}.$$

(ii) The GW energy loss scales like

$$\dot{E}_{\text{GW}} \sim \mathcal{M}_c^{\frac{10}{3}} f^{\frac{10}{3}}.$$

(iii) The rate of change of frequency scales like

$$\dot{f} \sim \mathcal{M}_c^{\frac{5}{3}} f^{\frac{11}{3}}.$$

---

\*Electronic address: [j.gair@ed.ac.uk](mailto:j.gair@ed.ac.uk)

(iv) The Fourier transform of  $h(t)$  scales like

$$\tilde{h} \sim \frac{1}{D} \mathcal{M}_c^{\frac{5}{6}} f^{-\frac{7}{6}}.$$

(v) The characteristic strain scales like

$$h_c \sim \frac{1}{D} \mathcal{M}_c^{\frac{5}{6}} f^{-\frac{1}{6}}.$$

(vi) The energy density of a GW background generated by a population of these sources scales like

$$\Omega_{\text{GW}}(f) \sim \mathcal{M}_c^{\frac{5}{3}} f^{\frac{2}{3}} \int_0^\infty \frac{N(z)}{(1+z)^{\frac{1}{3}}} dz.$$

In the above  $f$  denotes the orbital frequency of the binary and  $D$  the distance to the source.

If you made good notes during the lecture and are confident in those arguments, feel free to skip this and move straight on to the next part.

(2) We now suppose that there is an additional process driving the evolution of the binaries, stellar hardening. This leads to an evolution of the semi-major axis,  $a$ , of the binary of the form

$$\frac{d}{dt} \left( \frac{1}{a} \right) = k \frac{\rho_*}{\sigma^3} \frac{m_2}{a}$$

where  $k$  is a numerical constant,  $\rho_*$  is the stellar density and  $\sigma$  is the velocity dispersion of the stars. Show that this equation implies the energy loss from stellar hardening scales like

$$\dot{E}_{\text{hard}} \propto \frac{\rho m_2 \mu}{\sigma^3} M^{\frac{2}{3}} f^{\frac{2}{3}}. \quad (1)$$

(3) Derive an expression for the corresponding GW background energy density,  $\Omega_{\text{GW}}(f)$ .

(4) Assuming that the sources are at a common redshift (or the population is dominated by sources at a particular redshift), show that the spectrum is a broken power-law and find the asymptotic slopes at low and high frequency.

(5) If a broken power law background were detected, what would it tell you about the relevant physical processes influencing the population? How is that information encoded in the background?

(6) \* Try varying the assumptions about the population, e.g., the masses of the binary components or the properties of the stellar system or the redshift distribution of the sources. Compute the background from such populations numerically and explore how these assumptions change the results qualitatively and quantitatively.

## B. Burst sources

We now consider a population of burst sources. We will represent these as sine-Gaussians with a four parameter waveform family

$$h(t) = \frac{A}{D} \cos(2\pi f_0 t) e^{-\frac{1}{2} Q t^2}.$$

(1) Find the average waveform power

$$\langle h^2 \rangle = \frac{1}{2T} \int_{-T}^T h^2(t) dt$$

and show that, for a reasonable choice of  $T$ , e.g.,  $T = 2/\sqrt{Q}$ , it scales like  $A^2/D^2$  as expected. Why is this a 'reasonable choice' for  $T$ ?

(2) Compute the Fourier transform of  $h(t)$ . Hence deduce the bandwidth of the source is  $\Delta f \sim \sqrt{Q}/\pi$ .

(3) Estimate the SNR that might be possible in a burst search using

$$\left(\frac{S}{N}\right)^2 \approx \frac{\langle h^2 \rangle}{\Delta f S_n(f)}.$$

You may assume white noise,  $S_n(f) = \sigma^2$ , for simplicity.

- (4) Compute the SNR that would be obtained if the source was found by matched filtering and compare it to the burst search SNR. Comment on your answer.
- (5) Compute the energy distribution for one of these bursts,  $dE/df$ .
- (6) Find an expression for the GW background energy density produced by a population of these bursts, assuming that  $A$  and  $Q$  are constant and the distribution of  $f_0$  is a power law  $dn/df_0 \propto f_0^\alpha$  with  $\alpha > -1$ .
- (7) Assuming the sources are at a common redshift, show that the asymptotic slope of the background is  $f^3$  at low frequency and  $f^{3+\alpha}$  at high frequency.
- (8) \* Use simulations on a computer to explore how things change under modifications of these various assumptions.

### III. PART II: SAMPLING USING PYTHON

The second part of the workshop will be a computer practical. You will be sent a tarball containing a python notebook called `ASTROANDDAPRACTICAL.IPYNB`, plus some associated files. You will need to have jupyter botebook installed on your laptop and can then run this notebook using the command

```
jupyter-notebook AstroAndDAPractical.ipynb
```

There are a few packages that will be needed as well, specifically `CORNER`, `PYSTAN`, `SEABORN` and `H5PY`. These can be installed easily using `PIP`.

For the second part of the notebook you will need to download some LIGO public data. There are links in the notebook to the data, but the files are quite big.

